

**PATENT ABSTRACTS OF JAPAN**

(11)Publication number : 2002-034565

(43)Date of publication of application : 05.02.2002

---

(51)Int.Cl.	C12N 15/09
	A61K 38/00
	A61K 45/00
	A61P 31/04
	A61P 35/00
	A61P 37/08
	C07K 14/005
	C07K 16/28
	C12N 1/15
	C12N 1/19
	C12N 1/21
	C12N 5/10
	C12Q 1/68
	G01N 33/15
	G01N 33/50
	G01N 33/566
	G01N 33/577
	// C12P 21/08
	(C12Q 1/68
	C12R 1:91 )
	(C12P 21/08
	C12R 1:91 )

---

**BEST AVAILABLE COPY**

(21)Application number : 2000-219652

(71)Applicant : JAPAN SCIENCE & TECHNOLOGY  
CORP

(22)Date of filing : 19.07.2000

(72)Inventor : ARIRA SHIZUO  
HENMI HIROAKI

---

**(54) RECEPTOR PROTEIN SPECIFICALLY RECOGNIZING BACTERIAL DNA****(57)Abstract:**

**PROBLEM TO BE SOLVED:** To provide the receptor protein that specifically recognizes bacterial DNA bearing the non-methylated CpG sequence, the gene DNA coding the same and experimental model animals that are useful for examining the response of host immunocyte to bacterial infections.

**SOLUTION:** The DNA encoding the receptor protein that specifically recognizes the bacterial DNA bearing the non-methylated CpG sequence is screened by the BLAST search, a plurality of EST clones having high similarity to various kinds of TLS are screened and they are used as probes to isolate the full-length cDNA from the mouse macrophage cDNA library, the base sequence of the cDNA is analyzed to confirm that the TLR9 has the conserved areas such as the areas of LRR, TIR and the like. Then, the knockout mouse is established and the TLR9 is confirmed to be a receptor protein of the nucleotide including the non-methylated CpG sequence of the bacterial DNA.

---

**LEGAL STATUS**

[Date of request for examination] 09.08.2002

[Date of sending the examiner's decision of rejection] 09.03.2005

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision of rejection] 2005-06296

[Date of requesting appeal against examiner's decision of rejection] 08.04.2005

[Date of extinction of right]

Copyright (C); 1998,2003 Japan Patent Office

**\* NOTICES \***

JPO and NCIP I are not responsible for any damages caused by the use of this translation.

1.This document has been translated by computer. So the translation may not reflect the original precisely.

2.\*\*\*\* shows the word which can not be translated.

3.In the drawings, any words are not translated.

---

**CLAIMS**

---

[Claim(s)]

[Claim 1] DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array.

[Claim 2] DNA according to claim 1 to which receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array is characterized by being protein of the following (a) or (b).

(a) Protein which has reactivity to the bacterial DNA which 1 or some amino acid consist of deletion and an amino acid sequence permuted or added, and has a non-methylating CpG array in the amino acid sequence shown in the protein (b) array number 2 which consists of an amino acid sequence shown in the array number 2 [claim 3] DNA according to claim 1 characterized by including a part or all of these arrays in the base sequence shown in the array number 1, or its complementary sequence list.

[Claim 4] DNA according to claim 1 characterized by hybridizing under DNA which constitutes a gene according to claim 3, and stringent conditions.

[Claim 5] DNA according to claim 1 to which receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array is characterized by being protein of the following (a) or (b).

(a) Protein which has reactivity to the bacterial DNA which 1 or some amino acid consist of deletion and an amino acid sequence permuted or added, and has a non-methylating CpG array in the amino acid sequence shown in the protein (b) array number 4 which consists of an amino acid sequence shown in the array number 4 [claim 6] DNA according to claim 1 characterized by including a part or all of these arrays in the base sequence shown in the array number 3, or its complementary sequence list.

[Claim 7] DNA according to claim 1 characterized by hybridizing under DNA which constitutes a gene according to claim 6, and stringent conditions.

[Claim 8] Receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array.

[Claim 9] Protein according to claim 8 characterized by consisting of an amino acid sequence shown in the array number 2.

[Claim 10] Protein according to claim 8 characterized by 1 or some amino acid consisting of deletion and an amino acid sequence permuted or added in the amino acid sequence shown in the array number 2.

[Claim 11] Protein according to claim 8 characterized by consisting of an amino acid sequence shown in the array number 4.

[Claim 12] Protein according to claim 8 characterized by 1 or some amino acid consisting of deletion and an amino acid sequence permuted or added in the amino acid sequence shown in the array number 4.

[Claim 13] Claims 8-12 are fusion protein which combined protein, and the marker protein and/or the peptide tag of a publication either.

[Claim 14] Claims 8-12 are the antibodies specifically combined with the protein of a publication either.

[Claim 15] The antibody according to claim 14 characterized by an antibody being a monoclonal antibody.

[Claim 16] Claims 8-12 are the host cells which come to contain the manifestation system which can discover the protein of a publication either.

[Claim 17] The nonhuman animal characterized by the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array carrying out a superfluous manifestation.

[Claim 18] The nonhuman animal characterized by the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffering a loss on a chromosome.

[Claim 19] The nonhuman animal according to claim 18 characterized by being refractoriness to the bacterial DNA which has a non-methylating CpG array.

[Claim 20] Claims 17-19 to which the Rodentia animal is characterized by being a mouse are the nonhuman animals of a publication either.

[Claim 21] To the cell into which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, claims 1-7 are the preparation approaches of the cell which discovers the protein which has reactivity to the bacterial DNA which has the non-methylating CpG array characterized by introducing DNA of a publication either.

[Claim 22] The cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by being obtained by the preparation approach of the cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array according to claim 21.

[Claim 23] The screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by cultivating the cell which has discovered the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array by in vitro one under existence of a specimen material, and measuring and evaluating TLR9 activity, or an antagonist.

[Claim 24] The screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by to measure and evaluate the macrophage by which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array medicates with a specimen material the nonhuman animal which suffered a loss on the chromosome, and is obtained from this nonhuman animal, or the TLR9 activity of a spleen cell, or an antagonist.

[Claim 25] The screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by measuring and evaluating the macrophage by which the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array medicates with a specimen material the nonhuman animal which carried out the superfluous manifestation, and is obtained from this nonhuman animal, or the TLR9 activity of a spleen cell, or an antagonist.

[Claim 26] The screening approach of the agonist of the protein which has reactivity to the bacterial DNA which has the non-methylating CpG array according to claim 24 or 25 whose nonhuman animal is characterized by being a mouse, or an antagonist.

[Claim 27] either of claims 23-26 -- the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array acquired by the screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of a publication, or an antagonist, or an antagonist.

[Claim 28] The physic constituent which contains all of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, or its part as an active principle.

[Claim 29] The physic constituent which contains agonist or an antagonist according to claim 27 as an active principle.

[Claim 30] The diagnostic kit of the illness relevant to the deletion of the DNA array which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the

non-methylating CpG array characterized by including DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array in a specimen, and DNA according to claim 3 which can compare a base sequence with DNA according to claim 3, a permutation, and/or addition.

---

[Translation done.]

## \* NOTICES \*

JPO and NCIP are not responsible for any damages caused by the use of this translation.

1.This document has been translated by computer. So the translation may not reflect the original precisely.

2.\*\*\*\* shows the word which can not be translated.

3.In the drawings, any words are not translated.

---

## DETAILED DESCRIPTION

---

[Detailed Description of the Invention]

[0001]

[Field of the Invention] This invention relates to the genes of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, and this receptor protein, and those use.

[0002].

[Description of the Prior Art] It is known that a toll (Toll) gene is required for the decision (Cell 52, 269-279, 1988, Annu.Rev.Cell Dev.Biol.12, 393-416, 1996) of the dorso-ventral axis in the embryogenesis of drosophila and the antifungal immune response in an adult (Cell 86,973-983, 1996). This Toll is an I-beam film penetration acceptor which has a leucine rich repeat (LRR) to an extracellular field. This intracytoplasmic field It is clear that the intracytoplasmic field and homology of a mammals interleukin 1 acceptor (IL-1R) are high (). [ Nature 351, 355-356, 1991, ] [ Annu.Rev.Cell Dev.Biol.] 12, 393-416, 1996, J.Leukoc.Biol.63, 650-657, 1998.

[0003] The homologue of the mammals of Toll called a Toll Mr. acceptor (TLR) is identified in recent years. By current [, such as TLR2 and TLR4, ] Six families are reported (). [ Nature ] 388, 394-397, 1997, Proc.Natl.Acad.Sci.USA 95, 588-593, 1998, Blood 91, 4020-4027, 1998, Gene 231, 59-65, 1999. This TLR family minds MyD88 which is adapter protein like the above-mentioned IL-1R. Carry out the recruit of the IL-1R joint kinase (IRAK), and TRAF6 is activated. Activating down-stream NF-kappa B is known (J.Exp.Med.187, 2097-2101, 1998, Mol.Cell 2, 253-258, 1998, Immunity 11, 115-122, 1999). Moreover, it is thought that the role of the TLR family in the mammals is concerned with the immunological recognition native as a pattern recognition acceptor (PRR:pattern recognition receptor) which recognizes bacterial common structure (Cell 91, 295-298, 1997).

[0004] One of the pathogen associated-molecule patterns (PAMP:pathogen-associated molecular pattern) recognized by Above PRR It is lipopolysaccharide (LPS) which is the principal component of the adventitia of a gram negative (Cell 91, 295-298, 1997). This LPS stimulates a host cell. To a host cell TNFalpha, The various inflammatory cytokine of IL-1 and IL-6 grade is made to produce (457 28 Adv. Immunol. 293 -450, 1979, and Annu. Rev. Immunol. 13, 437- 1995), With LPS binding protein (LBP:LPS-bindingprotein) It is known that captured LPS will be handed over by CD14 on cell surface (Science 249, 1431-1433, 1990, Annu.Rev.Immunol.13, 437-457, 1995). It is unresponsive nature at LPS whose TLR4 knock AUTOUSU this invention persons produce the knockout mouse of TLR4, and is the principal component of the adventitia of the above-mentioned gram negative (J.Immunol.162, 3749-3752, and 1999), TLR2 knockout mouse was produced and it has reported that the reactivity over the peptidoglycan whose macrophages of TLR2 knockout mouse are a gram positive cell wall and its constituent falls (Immunity, 11, 443-451, and 1999).

[0005] On the other hand, an oligonucleotide including bacterial DNA (bacteria origin DNA) or a non-methylating CpG array The immunocyte of a mouse and Homo sapiens is stimulated (Trends Microbiol.4, 73-76, 1996, Trends Microbiol.6, 496-500, 1998), stimulating the T helper 1 cell (Th1) Mr. inflammatory response governed by emission of IL-12 and IFNgamma (EMBO J. -- 18, 6973-6982, 1999, J.Immunol.161, and 3042-3049 --) An oligonucleotide including a CpG array from 1998, Proc.Natl.Acad.Sci.USA 96, 9305-9310, and 1999 Cancer, As an adjuvant of the vaccine strategy containing the vaccine of allergy and an infectious disease \*\*\*\*\* is advocated

(Adv.Immunol.73, 329-368, 1999, Curr.Opin.Immunol.12, 35-43, 2000, Immunity 11, 123-129, 1999). Thus, in spite of expecting effectiveness in clinical practical use, the molecule mechanism in which bacterial DNA including a non-methylating CpG array activates immunocyte is not known well.

[0006]

[Problem(s) to be Solved by the Invention] As mentioned above, although the bacteria origin DNA containing the CpG motif which is not methylated activates immunocyte very much and the response of Th1 is guided, the activity with the molecular level is seldom understood. The technical problem of this invention is to offer a useful experiment model animal, when investigating the responsibility of the member receptor protein TLR9 of the TLR family which recognizes specifically the bacterial DNA which has a non-methylating CpG array which can clarify the operation with the molecular level of an oligonucleotide including the non-methylating CpG array of bacterial DNA, DNA which carries out the code of it, and the host immunocyte to a bacterial infectious disease.

[0007]

[Means for Solving the Problem] The TLR family in the mammals concerned with the immunological recognition native as a pattern recognition acceptor which recognizes bacterial common structure By current Six members (TLR 1-6) are released (Nature 388, 394-397, 1997, Proc.Natl.Acad.Sci.USA, 95, 588-593, 1998, Gene 231, 59-65, 1999). Two new members, TLR7 and TLR8, are registered into GenBank (registration numbers AF240467 and AF246971). moreover, the perfect length cDNA is found out also about TLR9, and it registers with GenBank -- \*\*\*\* (registration number AF 245704) -- it was not known about the function.

[0008] this invention persons screened DNA which carries out the code of the member receptor protein of the TLR family which recognizes specifically the bacterial DNA which has a non-methylating CpG array by BLAST search, screened many sequence tag (EST) clones which have the already identified various kinds TLR and high similarity, used such gene fragmentation as the probe, and isolated cDNA which has perfect die length from a mouse macrophage cDNA library, and Homo sapiens cDNA isolated them using this. Next, the base sequence of these cDNA(s) was analyzed and it checked that it was TLR9 to which saved areas, such as LRR and a TIR field, exist in this TLR family. Then, this TLR9 knockout mouse is produced, and it shows clearly that it is the receptor protein of an oligonucleotide with which TLR9 includes the non-methylating CpG array of bacterial DNA, and came to complete this invention.

[0009] Namely, DNA (claim 1) to which this invention carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, The receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array In the amino acid sequence shown in the protein (b) array number 2 which consists of an amino acid sequence shown in the DNA(a) array number 2 according to claim 1 characterized by being protein of the following (a) or (b) 1 or some amino acid consist of deletion and an amino acid sequence permuted or added. And the protein (claim 2) which has reactivity to the bacterial DNA which has a non-methylating CpG array, DNA (claim 3) according to claim 1 characterized by including a part or all of these arrays in the base sequence shown in the array number 1, or its complementary sequence list, DNA (claim 4) according to claim 1 characterized by hybridizing under DNA which constitutes a gene according to claim 3, and stringent conditions, The receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array In the amino acid sequence shown in the protein (b) array number 4 which consists of an amino acid sequence shown in the DNA(a) array number 4 according to claim 1 characterized by being protein of the following (a) or (b) 1 or some amino acid consist of deletion and an amino acid sequence permuted or added. And the protein (claim 5) which has reactivity to the bacterial DNA which has a non-methylating CpG array, DNA (claim 6) according to claim 1 characterized by including a part or all of these arrays in the base sequence shown in the array number 3, or its complementary sequence list, It is related with DNA (claim 7) according to claim 1 characterized by hybridizing under DNA which constitutes a gene according to claim 6, and stringent conditions.

[0010] Moreover, the receptor protein (claim 8) which recognizes specifically the bacterial DNA in which this invention has a non-methylating CpG array, In the protein (claim 9) according to claim 8 characterized by consisting of an amino acid sequence shown in the array number 2, and the amino acid sequence shown in the array number 2 The protein (claim 10) according to claim 8 characterized

by 1 or some amino acid consisting of deletion and an amino acid sequence permuted or added, In the protein (claim 11) according to claim 8 characterized by consisting of an amino acid sequence shown in the array number 4, and the amino acid sequence shown in the array number 4 It is related with the protein (claim 12) according to claim 8 characterized by 1 or some amino acid consisting of deletion and an amino acid sequence permuted or added.

[0011] Claims 8-12 this invention either Moreover, the fusion protein (claim 13) which combined protein, and the marker protein and/or the peptide tag of a publication, Claims 8-12 either The antibody (claim 14) specifically combined with the protein of a publication, It is related with the antibody (claim 15) according to claim 14 characterized by an antibody being a monoclonal antibody, and the host cell (claim 16) which comes to contain the manifestation system which can discover any of claims 8-12, or the protein of a publication.

[0012] Moreover, the nonhuman animal (claim 17) characterized by the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array carrying out the superfluous manifestation of this invention, The nonhuman animal (claim 18) characterized by the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffering a loss on a chromosome, It is related with the nonhuman animal (claim 19) according to claim 18 characterized by being refractoriness to the bacterial DNA which has a non-methylating CpG array, and the nonhuman animal (claim 20) of claims 17-19 to which the Rodentia animal is characterized by what is been a mouse which is a publication either.

[0013] This invention into moreover, the cell into which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome Claims 1-7 either The preparation approach (claim 21) of the cell which discovers the protein which has reactivity to the bacterial DNA which has the non-methylating CpG array characterized by introducing DNA of a publication, By the preparation approach of the cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array according to claim 21 It is related with the cell (claim 22) which discovers the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by being obtained.

[0014] Moreover, this invention cultivates the cell which has discovered the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array by in vitro one under existence of a specimen material. The screening approach (claim 23) of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by measuring and evaluating TLR9 activity, or an antagonist, The nonhuman animal to which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome is medicated with a specimen material. The screening approach (claim 24) of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by measuring and evaluating the macrophage obtained from this nonhuman animal, or the TLR9 activity of a spleen cell, or an antagonist, The nonhuman animal in which the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array carried out the superfluous manifestation is medicated with a specimen material. The screening approach (claim 25) of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by measuring and evaluating the macrophage obtained from this nonhuman animal, or the TLR9 activity of a spleen cell, or an antagonist, A nonhuman animal is related with the screening approach (claim 26) of the agonist of the protein which has reactivity to the bacterial DNA which has the non-methylating CpG array according to claim 24 or 25 characterized by being a mouse, or an antagonist.

[0015] Moreover, this invention Claims 23-26 either by the screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of a publication, or an antagonist The agonist or the antagonist (claim 27) of receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array acquired, The physic constituent (claim 28) which contains all of the receptor protein which recognizes



specifically the bacterial DNA which has a non-methylating CpG array, or its part as an active principle, DNA which carries out the code of the physic constituent (claim 29) which contains agonist or an antagonist according to claim 27 as an active principle, and the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array in a specimen, The deletion of the DNA array which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array characterized by including DNA according to claim 3 which can compare a base sequence with DNA according to claim 3, It is related with the diagnostic kit (claim 30) of the illness relevant to a permutation and/or addition.

[0016]

[Embodiment of the Invention] As bacterial DNA which has a non-methylating CpG array in this invention Immunocytes, such as a T cell, a B cell, and an antigen presenting cell, can be activated, and an immune response can be guided. As long as it is DNA originating in bacteria, such as oligodeoxynucleotide (ODN) which has the CpG motif which is not methylated, what kind of thing may be used. *ESERISHIA KORI*, the *Klebsiella pneumoniae*, *Pseudomonas AERUGINOSA*, *Salmonella typhimurium*, *Serratia marcescens*, the *Shigella flexneri*, *Vibrio KOREREE*, *Salmonella Minnesota*, *Porphyromonas gingivalis*, DNA of the bacteria origins, such as *Staphylococcus aureus*, the *Corynebacterium diphtheria*, *Nocardia KOERIAKA*, and *streptococcus new MONIA*, can be mentioned concretely.

[0017] As receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array It is not what will be restricted especially if it is protein which can recognize specifically the bacterial DNA which has a non-methylating CpG array. For example, it sets to the amino acid sequence shown by TLR9 and the array number 2 of the Homo sapiens origin shown by the array number 2 of an array table. The protein which can recognize specifically the bacterial DNA which 1 or some amino acid consist of deletion and an amino acid sequence permuted or added, and has the above-mentioned non-methylating CpG array, and these recombination protein can be mentioned concretely. The receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array can be prepared by the well-known approach based on the DNA array information etc.

[0018] moreover, as DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention In DNA which carries out the code of TLR9 of the Homo sapiens origin shown by the array number 2 of an array table, for example, the thing shown by the array number 1, and the amino acid sequence shown by the array number 2 1 or some amino acid consist of deletion and an amino acid sequence permuted or added. And DNA which carries out the code of the protein which can recognize specifically the bacterial DNA which has the above-mentioned non-methylating CpG array, DNA which carries out the code of the protein which can recognize specifically the bacterial DNA which hybridizes under these DNA and stringent conditions, and has the above-mentioned non-methylating CpG array is also included. These can be prepared by the well-known approach in TLR9 of the mouse origin based on the DNA array information etc. from a mouse RAW264.7cDNA library, 129 / SvJ mouse gene library, etc.

[0019] moreover, the thing for which the DNA which performs hybridization under stringent conditions to the DNA library of the mouse origin at the base sequence shown in the array number 1 or its complementary sequence list by using a part or all of these arrays as a probe, and hybridizes to this probe isolates -- receptor protein TLR9 -- said -- the DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the immunity induction non-methylating CpG array made into the \*\*\*\* purpose can also obtain. As conditions for the hybridization for acquiring this DNA, washing processing at 42 degrees C by 42-degree C hybridization and 1xSSC, and the buffer solution containing 0.1% of SDS can be mentioned, for example, and washing processing at 65 degrees C by 65-degree C hybridization and 0.1xSSC, and the buffer solution containing 0.1% of SDS can be mentioned more preferably. In addition, as an element which affects the stringency of hybridization, there are various elements in addition to the above-mentioned temperature conditions, and if it is this contractor, it is possible to realize stringency equivalent to the stringency of the hybridization which carried out [ above-mentioned ] instantiation, combining various elements suitably.

[0020] To the receptor protein which recognizes specifically the bacterial DNA which has non-methylating CpG arrays, such as a mouse and Homo sapiens, to be fusion protein of this invention What combined marker protein and/or a peptide tag is said. As marker protein As long as it is marker protein known conventionally, what kind of thing may be used. for example, can mention concretely alkaline phosphatase, Fc field of an antibody, HRP, GFP, etc., and as a peptide tag in this invention The peptide tag known conventionally [ , such as a Myc tag, a His tag, a FLAG tag and a GST tag, ] can be illustrated concretely. This fusion protein is useful also as purification of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array which could produce with the conventional method and used the compatibility of nickel-NTA and a His tag, detection of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, the quantum of an antibody to the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, and a reagent for research of this other these field.

[0021] As an antibody specifically combined with the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention A monoclonal antibody, a polyclonal antibody, a chimeric antibody, a single strand antibody, immunity, such as a hominization antibody, -- a specific antibody being mentioned concretely, and, although these can produce the bacterial DNA which has the above-mentioned non-methylating CpG array with a conventional method, using as an antigen the receptor protein recognized specifically A monoclonal antibody is more desirable in respect of the singularity also in it. a diagnosis of the illness to which the antibody specifically combined with the receptor protein which recognizes specifically the bacterial DNA which has non-methylating CpG arrays, such as this monoclonal antibody, originates in the variation or deletion of TLR9, and a controlled part of TLR9 -- a cordless handset -- it is useful when clarifying a style.

[0022] The antibody to the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array The fragment containing the receptor protein or the epitope which recognizes specifically the bacterial DNA which has this non-methylating CpG array to an animal (preferably except Homo sapiens) using the protocol of common use, It is produced by prescribing for the patient the cell which discovered this protein on the film front face. Or for preparation of a monoclonal antibody The hybridoma method for bringing about the antibody produced with the culture of continuation cell lineage (Nature 256, 495-497, 1975); The TORIOMA method, a Homo sapiens B cell hybridoma method (ImmunologyToday 4, 72, 1983) and the EBV-hybridoma method (MONOCLONAL ANTIBODIES AND CANCER THERAPY and pp.77-96 --) The approach of arbitration, such as Alan R.Liss, Inc., and 1985, can be used. As receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array below, the production approach of of the monoclonal antibody which mentions TLR9 of the mouse origin as an example, and is specifically combined to TLR9 of the mouse origin, i.e., anti-mTLR9 monoclonal antibody, is explained.

[0023] The above-mentioned anti-mTLR9 monoclonal antibody is producible by cultivating an anti-mTLR9 monoclonal-antibody production hybridoma with a conventional method by in vivo one or in vitro one. For example, in an in vivo system, it can obtain by cultivating by the culture medium for animal cell culture in an in vitro system a rodent and by cultivating by intraperitoneal [ of a mouse or a rat ] preferably again. As a culture medium for cultivating a hybridoma by the in vitro system, cell culture media, such as RPMI1640 containing antibiotics, such as streptomycin and penicillin, or MEM, can be illustrated.

[0024] An anti-mTLR9 monoclonal-antibody production hybridoma \*\*\*\* a BALB/c mouse using the receptor protein TLR9 obtained from the mouse etc., and can create an anti-mTLR9 monoclonal-antibody production hybridoma by carrying out the cell fusion of the spleen cell of a mouse and mouse NS-1 cell (ATCC TIB-18) by which immunity was carried out with a conventional method, and screening them with an immunofluorescent-stain pattern. Moreover, if it is the approach generally used to purification of protein as the separation / purification approach of this monoclonal antibody, what kind of approach may be used and liquid chromatography, such as affinity chromatography, can be illustrated concretely.

[0025] Moreover, in order to build the single strand antibody to the receptor protein which recognizes

specifically the bacterial DNA which has the above-mentioned non-methylating CpG array of this invention, the method of preparation (U.S. Pat. No. 4,946,778) of a single strand antibody is applicable. Moreover, in order to make a hominization antibody discover, a transgenic mouse or other mammals can be used, the clone which discovers the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array can be isolated and identified using the above-mentioned antibody, or affinity chromatography can also refine the polypeptide. The antibody to the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array is useful when clarifying the molecule device of receptor protein in which the bacterial DNA which has a non-methylating CpG array is recognized specifically. [0026] To antibodies, such as the above-mentioned anti-mTLR9 monoclonal antibody, moreover, for example, fluorescent materials, such as FITC (fluorescein isocyanate) or tetramethyl rhodamine isocyanate, The radioisotope of <sup>125</sup>I, <sup>32</sup>P, <sup>35</sup>S, or <sup>3</sup>H grade, and the alkaline phosphatase, By using the fusion protein with which what carried out the indicator with enzymes, such as a peroxidase, beta-galactosidase, or a phycoerythrin, firefly luminescence protein, such as Green fluorescence protein (GFP), etc. were united Functional analysis of the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array can be performed. Moreover, as an immunological measuring method, approaches, such as the RIA method, the ELISA method, a fluorescent antibody technique, a plaque technique, the spot method, an erythrocyte agglutination method, and an Ouchterlony method, can be mentioned.

[0027] This invention relates to the host cell which comes to contain the manifestation system which can discover the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array again. The installation to the host cell of the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array Davis et al. () [ BASIC METHODS ] IN MOLECULAR BIOLOGY, 1986, and Sambrook et al. () [ MOLECULAR ] CLONING: A LABORATORY the approach indicated by many standard laboratory manuals, such as MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., and 1989, -- for example Calcium phosphate transfection, DEAE-dextran medium transfection, Transformer BEKUSHON (transfection), a microinjection, Cationic lipid medium transfection, electroporation, transduction, scrape loading (scrape loading) Projectile installation (ballistic introduction), infection, etc. can perform. And as a host cell, animals-and-plants cells, such as fungus cells, such as bacteria prokaryotic cells, such as Escherichia coli, Streptomyces, a Bacillus subtilis, a streptococcus, and Staphylococcus, and yeast, an Aspergillus, the insect cell of Drosophila S2 and Spodoptera Sf9 grade, an L cell and a CHO cell, a COS cell, a HeLa cell, C127 cell, a BALB/c 3T3 cell (the variant which suffered a loss in dihydrofolate reductase, a thymidine kinase, etc. is included), a BHK-21 cell, HEK293 cell, a Bowes melanoma cell, and oocyte etc. can be mentioned.

[0028] Moreover, as long as it is the manifestation system which can make the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array as a manifestation system discover by host intracellular, what kind of thing may be used. The manifestation system originating in a chromosome, episome, and a virus, for example, the bacterial plasmid origin, The yeast plasmid origin, papovavirus like SV40, a vaccinia virus, Adenovirus, the fowlpox virus, the pseudorabies virus, the vector of the retrovirus origin, The vector originating in the bacteriophage origin, the transposon origins, and these combination, for example, the thing originating in the hereditary element of cosmid, a plasmid like phagemid, and a bacteriophage, can be mentioned. Not only making a manifestation cause but these manifestation system may include the control array which adjusts a manifestation.

[0029] The receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array which cultivates the cell membrane of the host cell which comes to contain the above-mentioned manifestation system, or this cell, and this cell, and is acquired can be used for the screening approach of this invention so that it may mention later. For example, as an approach of obtaining a cell membrane, F.Pietri-Rouxel's and others (Eur.J.Biochem., 247, 1174-1179, 1997) approach etc. can be used. Moreover, in order to collect and refine the receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array from a cell culture object An ammonium sulfate or ethanol precipitate, an acid extract, an anion, or a cation

exchange chromatography, A phosphocellulose chromatography, hydrophobic interaction chromatography, a well-known approach including affinity chromatography, hydroxyapatite chromatography, and a lectin chromatography -- high performance chromatography is used preferably. As a column especially used for affinity chromatography For example, the column which combined the receptor protein antibody which recognizes specifically the bacterial DNA which has anti-\*\* methylation CpG arrays, such as anti-TLR9 monoclonal antibody, When the usual peptide tag is added to the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of the above-mentioned TLR9 grade By using the column which combined the matter which has compatibility in this peptide tag, the receptor protein which recognizes specifically the bacterial DNA which has these non-methylating CpG arrays can be obtained.

[0030] With the nonhuman animal in which the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array in this invention carries out a superfluous manifestation The nonhuman animal which produces the receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array compared with a wild type nonhuman animal in large quantities is said. With moreover, the nonhuman animal by which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome A part or all of a gene that carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array on a chromosome is inactivated by gene variation, such as destruction, a deficit, and a permutation. A \*\*\*\*\* nonhuman animal is said for the function which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array. And as a nonhuman animal in this invention, although nonhuman animals, such as the Rodentia animals, such as a rabbit, and a mouse, a rat, can be mentioned concretely, it is not limited to these.

[0031] Moreover, refractoriness means the reactivity of the cell which constitutes the living body or living body to the stimulus by bacterial DNA, an organization, or an organ falling, or being lost mostly to the bacterial DNA which has a non-methylating CpG array in this invention. Therefore, the nonhuman animal of refractoriness means animals other than Homo sapiens, such as a mouse which the reactivity of the cell which constitutes a living body or a living body, an organization, or an organ is falling, or is lost mostly, a rat, and a rabbit, to the stimulus by bacterial DNA to the bacterial DNA which has a non-methylating CpG array in this invention. Moreover, as a stimulus by bacterial DNA, the in vivo stimulus which medicates a living body with bacterial DNA, the in vitro stimulus which contacts bacterial DNA into the cell separated from the living body can be mentioned, and, specifically, the nonhuman animal to which TLR9 gene functions, such as TLR9 knockout mouse, suffered a loss on the chromosome can be mentioned.

[0032] by the way, to the homozygote nonhuman animal born according to Mendel's laws The receptor protein deficit mold or superfluous manifestation mold which recognizes specifically the bacterial DNA which has a non-methylating CpG array, and the wild type of the brood are contained. From the ability of exact comparative experiments to be carried out on individual level by using the deficit mold or superfluous manifestation mold in a these homozygote nonhuman animal, and the wild type of the brood for coincidence The nonhuman animal of a wild type, i.e., the nonhuman animal to which the gene function which carries out a code suffers a loss or discovers [ superfluous ] the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array on a chromosome and an animal of the same kind, It is desirable to use the animal of a brood together on the occasion of screening of this invention indicated below furthermore. The gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array explains below the bacterial DNA which has a non-methylating CpG array for the production approach of the nonhuman animal which carries out a deficit or a superfluous manifestation taking the case of the knockout mouse and transgenic mouse of receptor protein which are recognized specifically on a chromosome.

[0033] For example, the mouse with which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of TLR9 grade suffered a loss on the chromosome, Namely, the receptor protein knockout mouse which recognizes specifically the bacterial DNA which has a non-methylating CpG array The

gene fragment obtained from the mouse gene library by approaches, such as PCR, is used. The gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array is screened. The subclone of the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the screened non-methylating CpG array is carried out using a virus vector etc., and it specifies by DNA sequencing. A target vector is produced by permuting all or a part of gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this clone by a pMC1 neo gene cassette etc., and introducing genes, such as a diphtheria toxin A fragmentation (DT-A) gene and a thymidine kinase (HSV-tk) gene of a herpes simplex virus, into 3' end side.

[0034] this produced targeting vector -- a line ---izing -- electroporation (electric punching) -- it introduces into an embryonic stem cell by law etc., homonous recombination is performed, and the embryonic stem cell which caused homonous recombination with antibiotics, such as G418 and gun cyclo beer (GANC), is chosen from those homonous recombinant. Moreover, it is desirable to check whether it is the recombinant which this selected embryonic stem cell makes the purpose with a Southern blot technique etc. The microinjection of the clone of the checked embryonic stem cell is carried out into the blastocyst of a mouse, this blastocyst is returned to the mouse of assumed parents, and a chimeric mouse is produced. If this chimeric mouse is made INTAKUROSU [ mouse / of a wild type ], the receptor protein knockout mouse which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention is producible by being able to obtain a heterozygote mouse and making this heterozygote mouse INTAKUROSU. Moreover, as an approach of checking whether the receptor protein knockout mouse which recognizes specifically the bacterial DNA which has a non-methylating CpG array having occurred, it investigates with a Northern blot technique etc., and there is a method of isolating RNA from the mouse obtained by the above-mentioned approach, and investigating the manifestation of this mouse by western blotting etc., for example.

[0035] That moreover, it is unresponsive nature to the bacterial DNA in which TLR9 created knockout mouse has a non-methylating CpG array For example, CpG ODN The macrophage of TLR9 knockout mouse, Immunocytes, such as a mononuclear cell and a dendritic cell, are made to contact by in vitro one or in vivo one. The amounts of production, such as TNF-alpha in this cell, IL-6, IL-12, and IFN-gamma, It can check by measuring activation of the molecule in the signal transfer path of TLR(s)9, such as the amount of manifestations of antigens, such as a growth response of a spleen B cell, and CD40, CD80 and CD86 in a spleen B cell front face, the MHC class II, and NF-kappa B and JNK, IRAK. And TLR9 knockout mouse of this invention can be used as a model useful to the vaccine development to an elucidation and bacterial infection of action mechanisms, such as bacterial DNA which has a non-methylating CpG array.

[0036] The transgenic mouse of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array The receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of TLR9 grade to cDNA which carries out a code A chicken beta-actin, Promotors, such as mouse neurofilament and SV40, and a rabbit beta globin, Unite poly A or the introns, such as SV40, and an introductory gene is built. The microinjection of this introductory gene is carried out to the pronucleus of a mouse fertilized egg. After cultivating the obtained ootid, it can transplant to the uterine tube of the mouse of assumed parents, and this transgenic mouse can be invented by breeding a transplanted animal after that and choosing the \*\* mouse which has said cDNA from the produced \*\* mouse. Moreover, selection of the \*\* mouse which has cDNA can be performed by the dot hybridization method which uses as a probe the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array which extracted and introduced rough DNA from the tail of a mouse etc., the PCR method using a specific primer, etc.

[0037] Moreover, if all or a part of DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention is used, a cell effective in gene therapies, such as deletion of receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, or an illness which originates unusually, can be prepared. As the preparation approach of these cells in this invention Into the cell

into which the gene function which carries out a code suffered a loss on the chromosome, the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array. All or a part of DNA of above-mentioned this invention is introduced with transfection etc. The approach of obtaining the cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array can be mentioned. It is desirable to use the cell which Above DNA etc. is integrated by the chromosome and shows TLR9 activity to a stay bull as a cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has this non-methylating CpG array especially.

[0038] And DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the above-mentioned non-methylating CpG array again, The antibody to the receptor protein which recognizes specifically the bacterial DNA which has the fusion non-methylating CpG array which combined the receptor protein, the marker protein, and/or the peptide tag which recognize specifically the bacterial DNA which has a non-methylating CpG array, The host cell which comes to contain the manifestation system which can discover the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, The nonhuman animal in which the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array carries out a superfluous manifestation, The nonhuman animal to which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, If the cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array is used The reactant inhibitor or reactant promoting agent to the agonist or the antagonist of receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention, and the bacterial DNA which has a non-methylating CpG array can be screened. In the inhibitor or promoting agent to the microbism, the inhibitor and preventive to an allergic disease or cancer, or a remedy, gene therapy, etc., it may be the matter useful for a diagnosis and therapies, such as an inhibitor or an inhibitor, and deletion of TLR9 activity or an illness which originates unusually, which was obtained by these screening about a side effect.

[0039] The TLR9 above-mentioned activity reacts specifically with the bacterial DNA which has a non-methylating CpG array, and means the function to transmit a signal to intracellular. As a signal communicative function The function which produces cytokine, such as TNF-alpha, IL-6, IL-12, and IFN-gamma, The function which produces nitrite ion, the function to increase a cell, and the function which discovers antigens, such as CD40, CD80, CD86, and the MHC class II, in cell surface, Although the function to activate the molecule in the signal transfer path of TLR(s)9, such as NF-kappa B and JNK and IRAK, etc. can be illustrated concretely, it is not limited to these.

[0040] As the screening approach of the agonist of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention, or an antagonist Immunocytes, such as the bottom of existence of a specimen material, a macrophage, a spleen cell, or a dendritic cell, The cell which has discovered the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, The cell which has discovered the protein which has reactivity to the bacterial DNA which has non-methylating CpG arrays, such as a cell which discovers the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, is cultivated by in vitro one. The approach of measuring and evaluating TLR9 activity, and a wild type nonhuman animal, the nonhuman animal, to which the gene function which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, Or the nonhuman animal in which the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array carried out the superfluous manifestation is medicated with a specimen material. The approach of measuring and evaluating the TLR9 activity of immunocytes, such as a macrophage obtained from this nonhuman animal, a spleen cell, or a dendritic cell, etc. can be mentioned concretely.

[0041] Moreover, since facing measuring and evaluating extent of macrophage activity or spleen cell activity, and comparing and estimating it as the measured value of a wild type nonhuman animal, especially the wild type nonhuman animal of a brood as contrast can abolish the variation by



individual difference, it is desirable. This is the same also in screening of the reactant inhibitor or promoting agent to the bacterial DNA which has the non-methylating CpG array shown below.

[0042] moreover, as the screening approach of of the reactant inhibitor or promoting agent to the bacterial DNA which has a non-methylating CpG array The protein which has reactivity to the bacterial DNA which has a non-methylating CpG array under existence with a specimen material and the bacterial DNA which has a non-methylating CpG array, Or the incubation of the cell membrane which has discovered this protein is carried out by in vitro one. The approach of measuring and evaluating reactivity with this protein, After making the macrophage or spleen cell obtained from the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, and a specimen material contact by in vitro one beforehand, This macrophage or a spleen cell is cultivated under existence of the bacterial DNA which has a non-methylating CpG array. The approach of measuring and evaluating extent of this macrophage, the macrophage activity of a spleen cell, or spleen cell activity, After making the macrophage or spleen cell obtained from the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, and the bacterial DNA which has a non-methylating CpG array contact by in vitro one beforehand, The approach of cultivating this macrophage or a spleen cell under existence of a specimen material, and measuring and evaluating extent of this macrophage, the macrophage activity of a spleen cell, or spleen cell activity, After medicating beforehand with a specimen material the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, The macrophage or spleen cell obtained from this nonhuman animal is cultivated under existence of the bacterial DNA which has a non-methylating CpG array. The approach of measuring and evaluating extent of this macrophage, the macrophage activity of a spleen cell, or spleen cell activity, After medicating beforehand with a specimen material the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, The approach of measuring and evaluating extent of the macrophage which this nonhuman animal is infected with bacteria and obtained from this nonhuman animal, the macrophage activity of a spleen cell, or spleen cell activity, After infecting beforehand the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome with bacteria, The macrophage or spleen cell obtained from this nonhuman animal is cultivated under existence of a specimen material. The approach of measuring and evaluating extent of this macrophage, the macrophage activity of a spleen cell, or spleen cell activity, After infecting beforehand the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome with bacteria, The approach of measuring and evaluating extent of the macrophage which medicates this nonhuman animal with a specimen material, and is obtained from this nonhuman animal, the macrophage activity of a spleen cell, or spleen cell activity, After medicating beforehand with a specimen material the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome, The approach of infecting this nonhuman animal with bacteria, and measuring and evaluating extent of the macrophage activity in this nonhuman animal, or spleen cell activity, After infecting beforehand the nonhuman animal to which the gene function which carries out the code of the protein which has reactivity to the bacterial DNA which has a non-methylating CpG array suffered a loss on the chromosome with bacteria, This nonhuman animal can be medicated with a specimen material, and the approach of measuring and evaluating extent of the macrophage activity in this nonhuman animal or spleen cell activity etc. can be mentioned concretely. Moreover, as bacterial DNA which has the non-methylating CpG array used for these screening approaches, it is CpG. Although it is desirable to use ODN (TCC-ATG-ACG-TTC-CTG-ATG-CT: array number 5), it is not limited to this, either.

[0043] The bacterial DNA which has the non-methylating CpG array which consists of comparing with the DNA array which carries out the code of the receptor protein which recognizes specifically the

bacterial DNA which has the non-methylating CpG array of this invention for the DNA array which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array in a specimen again is related in this invention to the diagnostic kit used for the diagnosis of the illness relevant to the activity of receptor protein or the manifestation recognized specifically. It is effective to the diagnosis of the illness produced by the too little manifestation of receptor protein which recognizes specifically the bacterial DNA which can perform detection of the variant of DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array by finding out the individual which has variation in a gene on DNA level, and has a non-methylating CpG array, a superfluous manifestation, or variation manifestation. Although the genomic DNA which can be obtained from biopsies, such as a test subject's cell, for example, blood, urine, saliva, and an organization, and RNA or cDNA can be concretely mentioned as a specimen used for this detection, when it is not limited to these and uses this specimen, what was amplified by PCR etc. can also be used. And change of the size of the magnification product when comparing with normal genotype can detect the deletion and insertion mutation of a base sequence, and point mutation can identify the receptor protein which recognizes specifically the bacterial DNA which has an indicator non-methylating CpG array for Magnification DNA by making it hybridize with the gene which carries out a code. Thus, the diagnosis or judgment of the illness relevant to the activity of receptor protein or the manifestation which recognizes specifically the bacterial DNA which has a non-methylating CpG array can be carried out by detecting the variation of the gene which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array.

[0044] This invention again A non-methylating CpG array The activity of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array which consists of all or a part of DNA which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which it has, or RNA of antisense strands, or the probe for a diagnosis of the disease relevant to a manifestation, And the antibody specifically combined with the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of the probe concerned and/or this invention is contained. It is related with the diagnostic kit of the disease relevant to the activity of receptor protein or the manifestation which recognizes specifically the bacterial DNA which has the becoming non-methylating CpG array. It is all or a part of DNA (cDNA) which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array as said probe for a diagnosis, or RNA (cRNA) of antisense strands, and it will not be restricted especially if it has the die length (at least 20 or more bases) of extent materialized as a probe. In order to make the antibody specifically combined with the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this probe and/or this invention into the active principle of the diagnostic drug of the disease of symptoms, such as microbism, it is desirable to dissolve in suitable buffers by which a probe is not disassembled, and sterilized water. Moreover, the immunity staining technique (Dev.Biol.170, 207-222, 1995, J. Neurobiol. 29, 1-17, 1996) using these diagnostic drugs, an In situ hybridization method (J. Neurobiol. 29, 1-17, 1996), and in situ The disease of symptoms, such as microbism, can also be diagnosed by approaches, such as the PCR method.

[0045] As long as all of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of TLR9 grade as a physic constituent of this invention or its part, the agonist of the above-mentioned receptor protein, and an antagonist are included, what kind of thing may be used. Specifically, the conquest agent, inhibitor, inhibitor, etc. of the side effect by existence of the CpG motif acting as a failure in the therapy and gene therapy using the vaccine to the microbism, the vaccine to cancer, the remedy of allergosis including bronchial asthma, and an antisense oligonucleotide can be mentioned.

[0046] As mentioned above, the deletion of the DNA array which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has the non-methylating CpG array of this invention, As a diagnostic kit of the illness relevant to a permutation and/or addition As long as DNA which carries out the code of TLR9 is included, what kind of thing may be used. By comparing a base sequence with DNA which carries out the code of the receptor protein which recognizes specifically DNA which carries out the code of this TLR9, and the bacterial DNA which has a



non-methylating CpG array in a specimen A diagnosis of the deletion of the DNA array which carries out the code of the receptor protein which recognizes specifically the bacterial DNA which has a non-methylating CpG array, a permutation and/or the illness relevant to addition, for example, cancer, allergy, an infectious disease, etc. is attained.

[0047]

[Example] Although an example is given to below and this invention is explained to it still more concretely, the technical range of this invention is not limited by these examples.

Example 1 (cloning of TLR9)

As a result of searching GenBank using Homo sapiens's TLR4 DNA array information, homology found out the very high mouse EST (registration number AA 273731; mouse). The mouse RAW264.7cDNA library was screened by having used the PCR magnification product of this mouse EST as the probe, and the cDNA clone of perfect length shown in the array number 3 containing TLR9 perfect open reading frame was isolated. GenBank was searched using the DNA array information on this mouse TLR9, and the human genome array which has high homology was found out. Based on this human genome array, the cDNA edge was amplified and cDNA of Homo sapiens TLR9 of the perfect length who has the base sequence shown in the array number 1 was isolated from U937 cell (J.Immunol.163, 5039-5048, 1999). [0048] Example 2 (production of TLR9 knockout mouse)

TLR9 genomic DNA was isolated from the 129-/SvJ mouse gene library (Stratagene make), the subclone was carried out in the pBluescript II SK(+) vector (Stratagene make), and it specified by restriction enzyme mapping and DNA sequencing. The targetting vector permuted the fragmentation of 1.0kbs which carry out the code of a part of LRR (leucine rich repeat) field by the neomycin resistance gene cassette (pMC1-neo; Stratagene make), and built it by inserting a herpes simplex virus thymidine kinase (HSV-TK) as a negative selective marker ( drawing 1 ). This targetting vector was line-ized, erection PORESHON was carried out, 292 clones which show resistance to G418 and gun cyclo beer were chosen as the day [ of viviparity / 14.1st ] embryonic stem cell (embryonic stem cell), and 14 clones were screened with the PCR method and a Southern blot technique.

[0049] The microinjection of the three target ES clones containing mutation TLR9 allele was carried out into the blastocyst of C57BL/6 mouse, and the chimeric mouse was produced. The chimeric mouse of this male was made to cross with C57BL / 6 female mouse, heterozygote F1 mouse was produced, and the homozygote mouse (TLR9 knockout mouse: TLR9-/-) was obtained by INTAKUROSU [ this heterozygote F1 mouse ] ( drawing 2 ). In addition, the check of a homozygote mouse digested each genomic DNA extracted from the tail of a mouse by ScaI, and was performed with the Southern blot technique using the probe shown in drawing 1 . TLR9 knockout mouse (TLR9-/-) of this invention could be produced according to Mendel's laws, and did not show remarkable abnormalities till the 12th week.

[0050] In order to check that inactivation of TLR9 gene has occurred by mutation, Apply to electrophoresis all RNA (10microg) extracted from the spleen cell of a wild type mouse (+/+) and TLR9 knockout mouse (-/-), and it moves to the nylon film. Northern blot analysis was performed to C-end fragmentation, N-end fragmentation, or beta-actin (beta-actin) of TLR9 which carried out the indicator using specific cDNA by [32P] ( drawing 3 R> 3). N-end fragmentation of these results to TLR9mRNA was not detected from the spleen cell of TLR9 knockout mouse. Moreover, when C-end fragmentation was used as a probe, although the thing of the wild type mouse origin and the thing of the almost same size were detected, in the volume, it turned out that the imprint of the mutant-mouse origin of Tlr9 is few. Then, RT-PCR method was performed using mRNA of the spleen cell obtained from the mutant mouse, and sequence analysis of the obtained product was performed. Consequently, it turned out that the neo gene is contained in Tlr9 imprinted gene, a stop codon appears at least in N-end of TLR9, and functional TLR9 protein is not discovered in a mutant mouse with this insertion of neo ( drawing 4 ). In addition, the extraordinary component was not seen as a result of measuring the lymph cell of TLR9 knockout mouse by flow cytometry.

[0051] Example 3 (preparation of a peritoneal macrophage)

It pours 2ml (product made from DIFCO) of 4% of thioglycollate media into intraperitoneal [ of a wild type mouse (wild-type) and TLR9 knockout mouse (TLR9-/-) / each ] at a time. Isolate a peritoneum exudate cell from intraperitoneal [ of each mouse ] three days after, and these cells are cultivated at 37 degrees C for 2 hours in the RPMI1640 culture medium (product made from GIBCO) which added

10% of foetal calf serum (product made from GIBCO). By washing with the Hanks buffer solution (product made from Hank's buffered salt solution:HBSS;GIBCO) of an ice temperature, the non-adherent cell was removed and it was used for the following experiments by making an adherent cell into a peritoneal macrophage.

[0052] Example 4 (responsibility over the bacterial DNA which has the non-methylating CpG array of TLR9 knockout mouse)

Recently, CpG It became clear to depend for the responsibility of ODN (oligodeoxynucleotide) on MyD88 which is adapter protein in the signal transfer path through TLR. This MyD88 knockout mouse is CpG. Although not answered to ODN, TLR2 knockout mouse and TLR4 knockout mouse are CpG normally. It answers to ODN. These things are CpG. It is shown that ODN is recognized by TLR(s) other than TLR2 and TLR4. Then, CpG of TLR9 knockout mouse The responsibility over ODN was investigated. First, the amount of production of the inflammatory cytokine in a peritoneal macrophage was measured as follows.

[0053] CpG of the various concentration shown under existence of INFgamma (30 unit/ml) or nonexistence at drawing 5 in each peritoneal macrophage prepared according to the example 3 It cultivated for 24 hours together with ODN (0.1 or the product made from 1.0microM;TIB MOLBIOL; TCC-ATG-ACG-TTC-CTG-ATG-CT), PGN (the product made from 10microg/ml;Sigma and Fluka; Staphylococcus-aureus origin), and LPS (the product made from 1.0microg/ml;Sigma; Salmonella Minnesota Re-595 origin). TNFalpha after culture and in a culture supernatant, IL-6, and IL-12 Each concentration of p40 was measured by the ELISA method. This result is shown in drawing 5. The macrophage of these results to a wild type mouse (Wild-type) is CpG. ODN is answered, TNFalpha, IL-6, and IL-12 are produced, and they are IFNgamma and CpG further. When stimulated by ODN, it turned out that the amount of production of TNFalpha, IL-6, and IL-12 increases. However, for the macrophage of the TLR9 knockout-mouse (TLR9-/-) origin, even the bottom of existence of IFNgamma is CpG. The inflammatory cytokine of detectable level was not produced in the response to ODN. Moreover, as for the macrophage of a wild type mouse and the TLR9 knockout-mouse origin, it turned out that comparable production of TNFalpha, IL-6, and IL-12 is mostly carried out by the response to LPS or PGN ( drawing 5 ). In addition, each experimental result shows the average of n= 3. It is shown that N.D. in drawing was undetectable.

[0054] Moreover, CpG It investigated about the responsibility of the spleen cell of the wild type mouse (Wild-type) to ODN or LPS, and TLR9 knockout mouse (TLR9-/-). CpG of the various concentration which isolates the spleen cell (1x10<sup>5</sup>) of each mouse, and is shown in drawing 6 It cultivated within 96 well plate by ODN or LPS, and the spleen cell was stimulated. 40 hours after culture, 1microcurie [3H]-thymidine (product made from DEYUPONTO) was added, it cultivated for further 8 hours, and the intake of [3H] was measured with beta scintillation counter (Packard make) ( drawing 6 ). At the spleen cell of this result to a wild type mouse, it is CpG. Although the cell proliferation reaction was promoted depending on the dose of ODN or LPS, at the spleen cell of TLR9 knockout mouse, it is CpG of what kind of concentration. It also sets to an ODN stimulus and is CpG. The cell proliferation reaction by ODN was not seen. Moreover, CpG ODN was answered and the manifestation of the major histocompatibility complex (MHC) class II on the front face of a B cell of the wild type mouse origin increased. However, at the B cell of the TLR9 knockout-mouse origin, it is CpG. The increment in the manifestation of the MHC class II guided to ODN was not seen. The macrophage and B cell of the above thing to TLR9 knockout mouse are CpG. It turned out that it is specifically lacking in the responsibility over ODN.

[0055] Next, CpG The bacteria origin DNA containing ODN stimulates a dendritic cell potentially, and supporting development of Th1 cell is known (EMBO J.18, 6973-6982, 1999, J.Immunol.161, 3042-3049, 1998, Proc.Natl.Acad.Sci.USA 96, 9305-9310, 1999). Then, CpG The up-regulation of the surface molecule of production of ODN induction cytokine and the dendritic cell of the bone marrow origin was analyzed. The bone marrow cell of a wild type mouse (Wild-type) or TLR9 knockout mouse (TLR9-/-) It cultivates by RPMI1640 culture medium which added 10% of foetal calf serum containing a 10 ng(s)/ml mouse granulocyte macrophage colony-stimulating factor (product made from Peprotech) (J.Exp.Med.176, 1693-1702, 1992). Immature dendritic cells will be collected after culture on the 6th, and it is CpG of 0.1microM. It cultivated for two days in the RPMI1640 culture medium which added 10% of foetal calf serum under existence of ODN or 0.1microg [/ml ] LPS or

nonexistence. IL-12 after culture and in supernatant liquid The concentration of p40 was measured by the ELISA method ( drawing 7 ). The dendritic cell of this result to the wild type mouse origin is CpG. Although ODN was answered and IL-12 were produced, it sets to the dendritic cell of the TLR9 knockout-mouse origin, and it is CpG. ODN did not guide production of IL-12.

[0056] It cultivates by RPMI1640 culture medium which added 10% of foetal calf serum containing a 10 ng/above-mentioned ml mouse granulocyte macrophage colony-stimulating factor (product made from Peprotech). In the dendritic cell collected on the 6th, receive CD40, CD80, CD86, and the MHC class II. Each biotin-ized antibody dyes and it is made to develop in the streptoavidin which carried out the indicator by the phycoerythrin (phycoerythrin:PE; product made from fur MINJIEN). Cel QUEST software (product made from BEKUTONDIKINSON) analyzed these cells by the fluorescence-activated-cell-sorter caliber (FACS Calibur) ( drawing 8 ). From this result to CpG Although the manifestation of CD40, CD80, CD86, and the MHC class II was promoted in the dendritic cell front face of the wild type mouse origin when stimulated by ODN, in the dendritic cell front face of the TLR9 knockout-mouse origin, it is CpG. The manifestation of these molecules was not promoted by the response to ODN ( drawing 8 ). In the stimulus by LPS, the response with the same said [ the dendritic cell of the wild type mouse origin ] of the dendritic cell of the TLR9 knockout-mouse origin was seen. The above result to TLR9 is CpG. It turned out that it is an acceptor indispensable to the cell response of ODN.

[0057] Example 5 (activation of NF-kappa B and JNK by the response to CpG ODN of the macrophage of the TLR9 knockout-mouse origin, and IRAK)

The signal of TLR activating IRAK which are a serine / threonine kinase through MyD88 which is an adapter molecule, and activating a MAP kinase and NF-kappa B subsequently is known (Immunity 11, 115-122, 1999). Then, CpG ODN investigated whether this intracellular signaling molecule would be activated. About the peritoneal macrophage (1x10<sup>6</sup>cells) of the wild type mouse prepared according to the example 3, and TLR9 knockout mouse, it is CpG of 1.0microM. The time amount stimulus was carried out, nucleoprotein was extracted from the macrophage of each mouse, it incubated together with the specific probe including the DNA bonding site of NF-kappa B shown in drawing 9 by ODN or LPS of 1.0microg [/ml ] Salmonella Minnesota Re-595, electrophoresis was performed, and it visualized with autoradiography ( drawing 9 R> 9).

[0058] From this result to CpG When stimulated by ODN, by the macrophage of the TLR9 knockout-mouse origin, the DNA avidity of NF-kappa B did not increase to the DNA avidity of NF-kappa B increasing in the macrophage of the wild type mouse origin. Activation of the NF-kappa B as what stimulated the macrophage of the wild type mouse origin by LPS with what [ same ] stimulated the macrophage of the TLR9 knockout-mouse origin by LPS was seen. From the above result to CpG It turns out that the activity of NF-kappa B by induction of ODN is specifically missing in the macrophage of the TLR9 knockout-mouse origin. In addition, the arrow head in drawing shows the location of the composite of NF-kappa B and a specific probe, and Yato shows the location of only a specific probe.

[0059] The time amount shown by drawing 10 or drawing 11 like the above, CpG The macrophage of the wild type mouse stimulated by ODN or LPS, and TLR9 knockout mouse Dissolution buffer solution (it NaCl(s) 1.0% of triton X-100 and 137mM by the last concentration) Tris of 20mM(s) - EDTA of HCl and 5mM, 10% of glycerol, PMSF of 1mM, the 20micro ag [/ml ] aprotinin, 20microg [/ml ] leupeptin, The buffer solution containing beta-glycerophosphoric acid of Na3VO4 of 1mM, and 10mM(s); It dissolves in pH8.0. Immunoprecipitation of this cell melt is carried out by the anti-JNK antibody (made in Santa Cruz), or the anti-IRAK antibody (wood primeval national-chemical-laborator incorporated company make). Like a reference (Immunity 11, 115-122, 1999) publication In vitro kinase assay was performed and the JNK activity which made the substrate GST-c-Jun dissolution protein (GST-c-Jun), and the activity of IRAK were measured ( drawing 10 , upper case;GST-c-Jun in 11, Auto).

[0060] Moreover, SDS-polyacrylamide gel electrophoresis was made to separate, the above-mentioned cell melt was moved to the nitrocellulose membrane, the blot of this film was carried out by the anti-JNK antibody (made in Santa Cruz), or the anti-IRAK antibody (product made from Transduction Laboratories), and it visualized using en HANSUDO chemiluminescence equipment (product made from DEYUPONTO) ( drawing 10 , the lower berth in 11; WB). From the above result

to CpG Although ODN activated JNK and IRAK of a macrophage of the wild type mouse origin, in the macrophage of the TLR9 knockout-mouse origin, it turned out that it is not activated at all ( drawing 10 , 11). Therefore, CpG It turned out that it depends for the signal transduction through ODN on TLR9.

[0061]

[Effect of the Invention] Although the bacteria origin DNA containing the CpG motif which is not methylated activated immunocyte very much and the response of Th1 was guided, the acceptor which recognizes the bacteria origin DNA was not known. Since the acceptor of an oligonucleotide including the non-methylating CpG array of bacterial DNA became clear by this invention, if the member receptor protein TLR9 of the TLR family which recognizes specifically the bacterial DNA which has a non-methylating CpG array, the gene DNA which carries out the code of it can be used for a diagnosis of a bacterially caused disease etc., and a therapy and a TLR9 knock-out animal is used, it will become possible [ clarifying the operation mechanism in the molecular level of the bacteria origin DNA ].

[0062]

[Layout Table]

SEQUENCE-LISTING <110> JAPAN SCIENCE AND-TECHNOLOGY-CORPORATION<120>

Specific-receptor-that-recognizes bacterial DNA<130> A031P63<140><141><160> 5 <170> PatentIn Ver. 2.1<210> 1<211> 3257<212> DNA<213> Homo sapiens<220> <221> CDS <222> .. (107) (3205)

<400> 1 ccgctgctgc ccctgtggga agggacctcg agtgtgaagc atccttcct gtagctgctg 60 tccagtctgc  
ccgccagacc ctctggagaa gcccctgccc cccagc atg ggt ttc 115 Met Gly Phe 1tgc cgc agc gcc ctg cac  
ccg ctg tct ctc ctg gtg cag gcc atc atg 163 Cys Arg Ser Ala Leu His Pro Leu Ser Leu Leu Val Gln  
Alalle Met 5 10 15 ctg gcc atg acc ctg gcc ctg ggt acc ttg cct gcc ttcta ccc tgt 211 Leu Ala Met  
Thr Leu Ala Leu Gly Thr Leu Pro Ala Phe Leu Pro Cys 20 25 30 35gag ctc cagccc cac ggc ctg gtg  
aac tgc aac tgg ctg ttc ctg aag 259 Glu Leu Gln Pro His Gly Leu Val Asn Cys Asn Trp Leu Phe Leu  
Lys 40 45 50 tct gtg ccc cacttc tcc atg gca gca ccc cgt ggc aat gtc acc agc 307 Ser Val Pro His Phe  
Ser Met Ala Ala Pro Arg Gly Asn Val Thr Ser 55 60 65 ctt tcc ttg tcc tcc aac cgc atc cac cac ctc  
cat gat tct gac ttt 355 Leu Ser Leu Ser Ser Asn Arg Ile His HisLeu His Asp Ser Asp Phe 70 75 80  
gcc cacctg ccc agc ctgogg cat ctc aac ctc aag tgg aac tgc ccg 403 Ala His Leu Pro Ser Leu Arg His  
Leu Asn Leu Lys Trp Asn Cys Pro 85 90 95 ccg gtt ggc ctc agc ccc atgcac ttc ccc tgc cac atg acc  
atc gag 451 Pro Val Gly Leu Ser Pro Met His Phe Pro Cys His Met Thr Ile Glu100 105 110 115ccc  
agc accttc ttg gct gtg ccc acc ctg gaa gag cta aac ctg agc 499 Pro Ser Thr Phe Leu Ala Val Pro Thr  
Leu Glu Glu Leu Asn Leu Ser 120 125 130 tac aac aac atc atg act gtg cct gcg ctg ccc aaa tcc ctc  
ata tcc 547 TyrAsn Asn Ile Met Thr Val Pro Ala Leu Pro Lys Ser Leu Ile Ser 135 140 145 ctg tcc ctc  
agc catacc aac atc ctg atg cta gac tct gcc agc ctc 595 Leu Ser Leu Ser His Thr Asn Ile Leu Met Leu  
Asp Ser Ala Ser Leu 150 155 160 gcc ggc ctg cat gcc-ctg-cgc-ttc-cta ttc-atg-gac-ggc-aac tgt tat  
643 Ala-Gly-Leu-His-Ala-Leu Arg Phe Leu Phe Met-Asp-Gly-Asn-Cys Tyr 165 170 175 tac aag aac  
ccc tgc-agg-cag-gca-ctg gag-gtg gcc ccg ggt gcc ctc 691 Tyr Lys Asn Pro Cys Arg Gln Ala Leu Glu  
Val Ala Pro Gly Ala Leu180 185 190 195ctt ggc ctg ggc aac ctc accacctg tca ctc aag tac aac aac  
ctc 739 Leu Gly Leu Gly Asn Leu Thr His Leu Ser Leu Lys Tyr Asn Asn Leu 200 205 210act gtg gtg  
ccc cgc aac ctg cct tcc agc ctg gag tat ctg ctg ttg 787 Thr Val Val Pro Arg Asn Leu Pro Ser Ser Leu  
Glu Tyr Leu Leu Leu 215 220 225 tcc tac aac cgc atcgtc aaa ctg gcg cct gag gac ctg gcc aat ctg 835  
Ser Tyr Asn Arg Ile Val Lys Leu Ala Pro Glu Asp Leu Ala Asn Leu 230 235 240 acc gcc ctg cgt gtg  
ctcgat gtg ggc gga aat tgc cgc cgc tgc gac 883 Thr Ala Leu ArgVal Leu Asp Val Gly Gly Asn Cys Arg  
Arg Cys Asp 245 250 255 cac gct ccc aac ccc tgc atggag tgc cct cgt cac ttc ccc cag cta 931 His Ala  
Pro Asn Pro Cys Met Glu Cys Pro Arg His Phe Pro Gln Leu260 265 270 275cat ccc gat acc ttc agc  
cac ctg agc cgt ctt gaa ggc ctg gtg ttg 979 His Pro Asp Thr Phe Ser His Leu Ser Arg Leu Glu Gly  
Leu Val Leu 280 285 290 aag gac agt tct ctc tcc tgg ctg aat gcc agt tgg ttc cgt ggg ctg 1027 Lys Asp  
Ser Ser Leu Ser Trp Leu Asn Ala Ser Trp Phe Arg Gly Leu 295 300 305 gga aac ctc cga gtg ctg gac  
ctg agt gag aac ttc ctc tac aaa tgc 1075 Gly Asn Leu Arg Val Leu Asp Leu Ser Glu Asn Phe Leu Tyr  
Lys Cys 310 315 320 atc act aaa acc aag gcc ttc cag ggc cta aca cag ctg cgc aag ctt 1123 Ile Thr  
Lys Thr Lys Ala Phe Gln Gly Leu Thr Gln Leu Arg Lys Leu 325 330 335 aac ctg tcc ttc aat tac caa  
aag agg gtg tcc ttt gcc cac ctg tct 1171 Asn Leu Ser Phe Asn Tyr Gln Lys Arg Val Ser Phe Ala His  
Leu Ser340 345 350 355ctg gcc-cct-tcc-ttc-ggg agc-ctg-gtc-gcc-ctg aag gag ctg gac atg 1219 Leu

Ala.Pro Ser Phe Gly Ser Leu-Val-Ala-Leu-Lys Glu Leu Asp Met 360 365 370 cac ggc atc ttc ttc cgc  
tca ctc gat gag acc acg ctc cgg-cca ctg 1267 His Gly Ile Phe Phe Arg Ser Leu Asp Glu Thr Thr Leu  
Arg Pro Leu 375 380 385 gcc cgc ctg ccc atg ctc cag act ctg cgt ctg cag atg aac ttc atc 1315 Ala  
Arg Leu Pro Met Leu Gln Thr Leu Arg Leu Gln Met Asn Phe Ile 390 395 400 aac cag gcc cag ctc ggc  
atc ttc agg gcc ttc cct ggc ctg cgc tac 1363 Asn Gln Ala Gln Leu Gly Ile Phe Arg Ala Phe Pro Gly  
Leu Arg Tyr 405 410 415 gtg gac ctg tcg gac aac cgc atc agc gga gct tog gag ctg aca gcc 1411 Val  
Asp Leu Ser Asp Asn Arg Ile Ser Gly Ala Ser Glu Leu Thr Ala420 425 430 435acc atg ggg gag gca gat  
ggaggggag aag gtc tgg ctg cag cct ggg 1459 Thr Met Gly Glu Ala Asp Gly Gly Glu Lys Val Trp Leu Gln  
Pro Gly 440 445 450 gac ctt gct ccg gcc cca gtg gac act ccc agc tct gaa gac ttc agg 1507 Asp Leu  
Ala Pro Ala Pro Val Asp Thr Pro Ser Ser Glu Asp Phe Arg 455 460 465 ccc aac tgc agc acc ctc aac  
ttc acc ttg gat ctg tca cgg aac aac 1555 Pro Asn Cys Ser Thr Leu Asn Phe Thr Leu Asp Leu Ser Arg  
Asn Asn 470 475 480 ctg gtg acc gtg cag ccg gag atg ttt gcc cag ctc tcg cac ctg cag 1603 Leu Val  
Thr Val Gln Pro Glu Met Phe Ala Gln Leu Ser His Leu Gln 485 490 495 tgc ctg cgc ctg agc cac aac  
tgc atc tcg cag gca gtc aat ggc tcc 1651 Cys Leu Arg Leu Ser His Asn Cys Ile Ser Gln Ala Val Asn  
Gly Ser500 505 510 515cag ttc ctg ccg ctg acc ggtctgcag gtg cta gac ctg tcc cac aat 1699 Gln Phe  
Leu Pro Leu Thr Gly Leu Gln Val Leu Asp Leu Ser His Asn 520 525 530 aag ctg gac ctc tac ca  
c-gag-cac-tca-ttc-acg gag cta cca cga ctg 1747 Lys Leu Asp Leu Tyr His  
Glu-His-Ser-Phe-Thr-Glu-Leu-Pro-Arg-Leu 535 540 545 gag gcc ctg gac ctc agc tac aac agc cag  
ccc ttt ggc atg cag ggc 1795 Glu Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe Gly Met Gln Gly 550  
555 560 gtg ggc cac aac ttc agc ttc gtg gct cac ctg cgc acc ctg cgc cac 1843 Val Gly His Asn Phe  
Ser Phe Val Ala His Leu Arg Thr Leu Arg His 565 570 575 ctc agc ctg gcc cac aac aac atc cac agc  
caa gtg tcc cag cag ctc 1891 Leu Ser Leu Ala His Asn Asn Ile His Ser Gln Val Ser Gln Gln Leu580  
585 590 595tgc agt acg tcg ctg cgg gcctggac ttc agc ggc aat gca ctg ggc 1939 Cys Ser Thr Ser Leu  
Arg Ala Leu Asp Phe Ser Gly Asn Ala Leu Gly 600 605 610 cat atg tgg gcc gag gga gac ctc tat ctg  
cac ttc ttc caa ggc ctg 1987 His Met Trp Ala Glu Gly Asp Leu Tyr Leu His Phe Phe Gln Gly Leu 615  
620 625 agc ggt ttg atc tgg ctg gac ttg tcc cag aac cgc ctg cac acc ctc 2035 Ser Gly Leu Ile Trp Leu  
Asp Leu Ser Gln Asn Arg Leu His Thr Leu 630 635 640 ctg ccc caa acc ctg cgc aac ctc ccc aag agc  
cta cag gtg ctg cgt 2083 Leu Pro Gln Thr Leu Arg Asn Leu Pro Lys Ser Leu Gln Val Leu Arg 645 650  
655 ctc cgt gac aat tac ctg gcc ttc ttt aag tgg tgg agc ctc cac ttc 2131 Leu Arg Asp Asn Tyr Leu Ala  
Phe Phe Lys Trp Trp Ser Leu His Phe660 665 670 675ctg ccc aaa ctg gaa gtc ctogacctg gca gga aac  
cag ctg aag gcc 2179 Leu Pro Lys Leu Glu Val Leu Asp Leu Ala Gly Asn Gln Leu Lys Ala 680 685  
690 ctg acc aat ggc agc ctg cct gct ggc acc cgg ctc cgg agg ctg gat 2227 Leu Thr Asn Gly Ser Leu  
Pro Ala Gly Thr Arg Leu Arg Arg Leu Asp 695 700 705 gtc agc tgc aac agc atc agc ttc gtg gcc ccc  
ggc ttc ttt tcc aag 2275 Val Ser Cys Asn Ser Ile Ser Phe Val Ala Pro Gly Phe Phe Ser Lys 710 715  
720 gcc aag gag ctg cga gag ctc aac ctt agc gcc aac gcc ctc aag aca 2323 Ala Lys Glu Leu Arg Glu  
Leu Asn Leu Ser Ala Asn Ala Leu Lys Thr 725 730 735 gtg gac cac tcc tgg ttt ggg ccc ctg gcg agt  
gcc ctg caa ata cta 2371 Val Asp His Ser Trp Phe-Gly-Pro-Leu-Ala Ser Ala Leu Gln Ile-Leu740 745  
750 755gat gta agc gcc aac cct-ctg-cac-tgc-gcc tgt ggg gcg gcc ttt atg 2419 Asp Val Ser Ala Asn  
Pro Leu-His-Cys-Ala-Cys Gly Ala Ala Phe Met 760 765 770 gac ttc ctg ctg gaggtgcag gct gcc gtg  
ccc ggt ctg ccc agc cgg 2467 Asp Phe Leu Leu Glu Val Gln Ala Ala Val Pro Gly Leu Pro Ser Arg 775  
780 785 gtg aag tgt ggc agtccg ggc cag ctc cag ggc ctc agc atc ttt gca 2515 Val Lys Cys Gly Ser Pro  
Gly Gln Leu Gln Gly Leu Ser Ile Phe Ala 790 795 800 cag gacctg cgc ctc tgcctg gat gag gcc ctc tcc  
tgg gac tgt ttc 2563 Gln Asp Leu Arg Leu Cys Leu Asp Glu Ala Leu Ser Trp Asp Cys Phe 805 810  
815 gcc ctc tcg ctg ctg gct gtggct ctg ggc ctg ggt gtg ccc atg ctg 2611 Ala Leu Ser Leu Leu Ala Val  
Ala Leu Gly Leu Gly Val Pro Met Leu820 825 830 835cat cac ctc tgt ggc tgg gac ctc tgg tac tgc ttc  
cac ctg tgc ctg 2659 His His Leu Cys Gly Trp Asp Leu Trp Tyr Cys Phe His Leu Cys Leu 840 845  
850 gcc tgg ctt ccc tgg cgg ggg cgg caa agt ggg cga gat gag gat gcc 2707 Ala Trp Leu Pro Trp Arg  
Gly Arg Gln Ser Gly Arg Asp Glu Asp Ala 855 860 865 ctg ccc tac gat gccttc gtg gtc ttc gac aaa acg  
cag agcgca gtg 2755 Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Thr Gln Ser Ala Val 870 875 880  
gca gac tgg gtg tac aacgag ctt cgg ggg cag ctg gag gag tgc cgt 2803 Ala Asp Trp Val Tyr Asn Glu Leu  
Arg Gly Gln Leu Glu Glu Cys Arg 885 890 895 ggg cgc tgg gcactc cgc ctgtgc ctggag gaa cgc gac tgg  
ctg cct 2851 Gly Arg Trp Ala Leu Arg Leu Cys Leu Glu Glu Arg Asp Trp Leu Pro900 905 910 915ggc  
aaa acc ctc ttt gag aac ctg tgg gcc tcg gtc tat ggc agc cgc 2899 Gly Lys Thr Leu Phe Glu Asn Leu  
Trp Ala Ser Val Tyr Gly Ser Arg 920 925 930 aag acg ctg ttt gtg ctg gcc cac acg gac cgg gtc agt ggt

ctc ttg 2947 Lys Thr Leu Phe Val Leu Ala His Thr Asp Arg Val Ser Gly Leu Leu 935 940 945 cgc gcc  
agc ttc ctg-ctg-gcc-cag-cag cgc ctg ctg gag gac cgc aag 2995 Arg Ala Ser Phe Leu  
Leu-Ala-Gln-Gln-Arg Leu Leu Glu Asp Arg-Lys 950 955 960 gac gtc gtg gtg ctg gtg atc ctg agc cct  
gac ggc cgc cgc tcc cgc 3043 Asp Val Val Val Leu Val Ile Leu Ser Pro Asp Gly Arg Arg Ser Arg 965  
970 975 tac gtg cgg ctg cgc cag cgc ctc tgc cgc cag agt gtc ctc ctc tgg 3091 Tyr Val Arg Leu Arg  
Gln Arg Leu Cys Arg Gln Ser Val Leu Leu Trp980 985 990 995ccc cac cag ccc agt ggt cagcgcagc ttc  
tgg gcc cag ctg ggc atg 3139 Pro His Gln Pro Ser Gly Gln Arg Ser Phe Trp Ala Gln Leu Gly Met 1000  
1005 1010 gcc ctg acc agg gac aac cac cac ttc tat aac cgg aac ttc tgc cag 3187 Ala Leu Thr Arg  
Asp Asn His His Phe Tyr Asn Arg Asn Phe Cys Gln 1015 1020 1025 gga ccc acg gcc gaa tag  
ccgtgagccg gaatctgca cgggtgccacc 3235 Gly Pro Thr Ala Glu 1030 tccacactca cctcacctct gc 3257  
<210> 2 <211> 1032<212> PRT <213> Homo sapiens <400> 2 Met Gly Phe Cys Arg Ser Ala Leu His  
Pro Leu Ser Leu Leu Val Gln 1 5 10 15 Ala Ile Met Leu Ala Met Thr Leu Ala LeuGly Thr Leu Pro Ala  
Phe 20 25 30 Leu Pro Cys Glu Leu Gln Pro His Gly Leu Val Asn Cys Asn Trp Leu 35 40 45 Phe Leu  
Lys Ser Val Pro His Phe Ser Met Ala Ala Pro Arg Gly Asn 50 55 60 Val Thr SerLeu SerLeu Ser Ser  
Asn Arg Ile His His Leu His Asp 65 70 75 80 Ser Asp Phe Ala His Leu Pro Ser Leu Arg His Leu Asn  
Leu Lys Trp 85 90 95 Asn Cys Pro Pro Val Gly Leu Ser Pro Met His Phe Pro Cys His Met 100 105  
110 Thr Ile Glu Pro Ser Thr Phe Leu AlaVal Pro Thr Leu Glu Glu Leu 115 120 125 Asn Leu Ser Tyr  
Asn Asn Ile Met Thr-Val-Pro-Ala-Leu-Pro-Lys-Ser 130 135 140 Leu Ile Ser Leu Ser Leu Ser His  
Thr-Asn-Ile-Leu-Met-Leu Asp Ser 145 150 155 160 Ala Ser Leu Ala Gly Leu His Ala Leu Arg Phe  
Leu Phe Met Asp Gly 165 170 175 Asn Cys Tyr Tyr Lys Asn Pro Cys Arg Gln Ala Leu Glu Val Ala  
Pro 180 185 190 Gly Ala Leu Leu Gly Leu Gly Asn Leu Thr His Leu Ser Leu Lys Tyr 195 200 205 Asn  
Asn Leu Thr Val Val Pro Arg Asn Leu Pro Ser Ser Leu Glu Tyr 210 215 220 Leu Leu Leu Ser Tyr  
Asn Arg Ile Val Lys Leu Ala Pro Glu Asp Leu 225 230 235 240 Ala Asn Leu Thr Ala Leu Arg Val Leu  
Asp Val Gly Gly Asn Cys Arg 245 250 255 Arg Cys Asp His Ala Pro Asn Pro Cys Met Glu Cys Pro  
Arg His Phe 260 265 270 Pro Gln Leu His Pro Asp Thr Phe Ser His Leu Ser Arg Leu Glu Gly 275 280  
285 Leu Val Leu Lys Asp Ser Ser Leu Ser Trp Leu Asn Ala Ser Trp Phe 290 295 300 Arg Gly Leu  
Gly Asn Leu Arg Val Leu Asp Leu Ser Glu Asn Phe Leu 305 310 315 320 Tyr Lys Cys Ile Thr Lys Thr  
Lys Ala Phe Gln Gly Leu Thr Gln Leu 325 330 335 Arg Lys Leu Asn Leu Ser Phe Asn Tyr Gln Lys  
Arg Val Ser Phe Ala 340 345 350 His Leu Ser Leu Ala Pro Ser Phe Gly Ser Leu Val Ala Leu Lys Glu  
355 360 365 Leu Asp Met His Gly Ile Phe Phe Arg Ser Leu Asp Glu Thr Thr Leu 370 375 380 Arg Pro  
Leu Ala Arg Leu Pro Met Leu Gln Thr Leu Arg Leu Gln Met 385 390 395 400 Asn Phe Ile Asn Gln Ala  
Gln Leu Gly Ile Phe Arg Ala Phe Pro Gly 405 410 415 Leu Arg Tyr Val Asp Leu Ser Asp Asn Arg Ile  
Ser Gly Ala Ser Glu 420 425 430 Leu Thr Ala Thr Met Gly Glu Ala Asp Gly Gly Glu Lys Val Trp Leu  
435 440 445 Gln Pro Gly Asp Leu Ala Pro Ala Pro Val Asp Thr Pro Ser Ser Glu 450 455 460 Asp Phe  
Arg Pro Asn Cys Ser Thr Leu-Asn-Phe-Thr-Leu-Asp Leu Ser 465 470 475 480 Arg Asn Asn Leu Val  
Thr Val Gln Pro-Glu-Met-Phe-Ala-Gln-Leu-Ser 485490 495 His Leu Gln Cys Leu Arg Leu Ser His  
Asn Cys Ile Ser Gln Ala Val 500 505 510 Asn Gly Ser Gln Phe Leu Pro Leu Thr Gly Leu Gln Val Leu  
Asp Leu 515 520 525 Ser His Asn Lys Leu Asp Leu Tyr His Glu His Ser Phe Thr Glu Leu 530 535  
540 Pro Arg Leu Glu Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe Gly 545 550 555 560 Met Gln  
Gly Val Gly His Asn Phe Ser Phe Val Ala His Leu Arg Thr 565 570 575 Leu Arg His Leu Ser Leu Ala  
His Asn Asn Ile His Ser Gln Val Ser 580 585 590 Gln Gln Leu Cys Ser Thr Ser Leu Arg Ala Leu Asp  
Phe Ser Gly Asn 595 600 605 Ala Leu Gly His Met Trp Ala Glu Gly Asp Leu Tyr Leu His Phe Phe 610  
615 620 Gln Gly Leu Ser Gly Leu Ile Trp Leu Asp Leu Ser Gln Asn Arg Leu 625 630 635 640 His Thr  
Leu Leu Pro Gln Thr Leu Arg Asn Leu Pro Lys Ser Leu Gln 645 650 655 Val Leu Arg Leu Arg Asp  
Asn Tyr Leu Ala Phe Phe Lys Trp Trp Ser 660 665 670 Leu His Phe Leu Pro Lys Leu Glu Val Leu  
Asp Leu Ala Gly Asn Gln 675 680 685 Leu Lys Ala Leu Thr Asn Gly Ser Leu Pro Ala Gly Thr Arg Leu  
Arg 690 695 700 Arg Leu Asp Val Ser Cys Asn Ser Ile Ser Phe Val Ala Pro Gly Phe 705 710 715 720  
Phe Ser Lys Ala Lys Glu Leu Arg Glu Leu Asn Leu Ser Ala Asn Ala 725 730 735 Leu Lys Thr Val Asp  
His Ser Trp Phe Gly Pro Leu Ala Ser Ala Leu 740 745 750 Gln Ile Leu Asp Val Ser Ala Asn Pro Leu  
His Cys Ala Cys Gly Ala 755 760 765 Ala Phe Met Asp Phe Leu Leu Glu Val Gln Ala Ala Val Pro Gly  
Leu 770 775 780 Pro Ser Arg Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Leu Ser 785 790 795 800  
Ile Phe Ala Gln Asp Leu Arg Leu Cys-Leu-Asp-Glu-Ala-Leu-Ser-Trp 805 810 815 Asp Cys Phe Ala  
Leu Ser Leu Leu Ala-Val-Ala-Leu-Gly Leu Gly Val 820 825830 Pro Met Leu His His Leu Cys Gly Trp  
Asp Leu Trp Tyr Cys Phe His 835 840 845 Leu Cys Leu Ala Trp Leu Pro Trp Arg Gly Arg Gln Ser Gly



Arg Asp 850 855 860 Glu Asp Ala Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Thr Gln 865 870 875  
 880 Ser Ala Val Ala Asp Trp Val Tyr Asn Glu Leu Arg Gly Gln Leu Glu 885 890 895 Glu Cys Arg Gly  
 Arg Trp Ala Leu Arg Leu Cys Leu Glu Glu Arg Asp 900 905 910 Trp Leu Pro Gly Lys Thr Leu Phe  
 Glu Asn Leu Trp Ala Ser Val Tyr 915 920 925 Gly Ser Arg Lys Thr Leu Phe Val Leu Ala His Thr Asp  
 Arg Val Ser 930 935 940 Gly Leu Leu Arg Ala Ser Phe Leu Leu Ala Gln Gln Arg Leu Leu Glu 945 950  
 955 960 Asp Arg Lys Asp Val Val Val Leu Val Ile Leu Ser Pro Asp Gly Arg 965 970 975 Arg Ser Arg  
 Tyr Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val 980 985 990 Leu Leu Trp Pro His Gln Pro Ser  
 Gly Gln Arg Ser Phe Trp Ala Gln 995 1000 1005 Leu Gly Met AlaLeu Thr Arg Asp Asn His His Phe  
 Tyr Asn Arg Asn 1010 1015 1020 Phe Cys Gln GlyProThr Ala Glu 1025 1030 <210> 3 <211>  
 3471<212> DNA <213> Mus musculus <220> <221> CDS <222> .. (107) (3205) <400> 3 tgaaagtgtc  
 acttcctcaa ttctctgaga gaccctgggtg tggaacatca ttctctgcgc 60 cccagtttgt cagagggagc ctccggagaa  
 tcctccatct cccaac atg gtt ctc 115 Met Val Leu 1cgt cga agg act ctg cac-ccc-ttg-tcc-ctc ctg gta cag  
 gct gca gtg 163 Arg Arg Arg Thr Leu His Pro-Leu-Ser-Leu-Leu Val Gln Ala Ala Val 5 10 15 ctg gct  
 gag act ctg gcc ctg ggt acc ctg cct gcc ttc cta ccc tgt 211 Leu Ala Glu ThrLeu Ala Leu Gly Thr Leu  
 Pro Ala Phe Leu Pro Cys 20 25 30 35gag ctg aagcct cat ggc ctg gtggac tgc aat tgg ctg ttc ctg aag  
 259 GluLeu Lys Pro His Gly Leu Val Asp Cys Asn Trp Leu Phe Leu Lys 40 45 50 tct gta ccc cgtttc  
 tct gcg gca gca tcc tgc tcc aac atc acc cgc 307 Ser Val Pro Arg Phe Ser Ala Ala Ala Ser Cys Ser  
 Asn Ile Thr Arg 55 60 65 ctc tcc ttg atc tccaac cgt atc cac cac ctg cac aac tcc gac ttc 355 Leu Ser  
 Leu Ile Ser Asn Arg Ile His His Leu His Asn Ser Asp Phe 70 75 80 gtc cac ctg tcc aac ctgcgg cag ctg  
 aac ctc aag tgg aac tgt cca 403 Val His Leu SerAsn Leu Arg Gln Leu Asn Leu Lys Trp Asn Cys Pro  
 85 90 95 ccc act ggc ctt agc ccc ttgcac ttc tct tgc cac atg acc att gag 451 Pro Thr Gly Leu Ser Pro  
 Leu His Phe Ser Cys His Met Thr Ile Glu100 105 110 115ccc aga accttc ctg gct atg cgt aca ctg gag  
 gag ctg aac ctg agc 499 Pro Arg Thr Phe Leu Ala Met Arg Thr Leu Glu Glu Leu Asn Leu Ser 120 125  
 130 tat aat ggt atc acc act gtg ccc cga ctg ccc agc tcc ctg gtg aat 547 TyrAsn Gly Ile Thr Thr Val  
 Pro Arg Leu Pro Ser Ser Leu Val Asn 135 140 145 ctg agc ctg agc cacacc aac atc ctg gtt cta gat gct  
 aac agc ctc 595 Leu Ser Leu Ser His Thr Asn Ile Leu Val Leu Asp Ala Asn Ser Leu 150 155 160 gcc  
 ggcta tac agc ctgcgc gtt ctc ttc atg gac ggg aac tgc tac 643 Ala Gly Leu Tyr Ser Leu Arg Val Leu  
 Phe Met Asp Gly Asn Cys Tyr 165 170 175 tac aag aac ccc tgc aca ggagcg gtg aag gtg acc cca ggc  
 gcc ctc 691 Tyr Lys Asn Pro Cys Thr Gly Ala Val Lys Val Thr Pro Gly Ala Leu180 185 190 195ctg  
 ggc ctg agc aat ctc acc cat ctg tct gtg aag tat aac aac ctc 739 Leu Gly Leu Ser Asn Leu Thr His Leu  
 Ser Val Lys Tyr Asn Asn Leu 200 205 210 aca aag gtg ccc cgc-caa-ctg-ccc-ccc agc ctg gag tac ctc  
 ctg gtg 787 Thr Lys Val Pro Arg Gln-Leu-Pro-Pro-Ser Leu Glu Tyr Leu Leu-Val 215 220 225 tcc tat  
 aac ctc att-gtc-aag-ctg-ggg cct gaa gac ctg gcc aat ctg 835 Ser Tyr Asn Leu Ile Val Lys Leu Gly  
 Pro Glu Asp Leu Ala Asn Leu 230 235 240 acc tcc ctt cga gta ctt gat gtg ggt ggg aat tgc cgt cgc tgc  
 gac 883 Thr Ser Leu Arg Val Leu Asp Val Gly Gly Asn Cys Arg Arg Cys Asp 245 250 255 cat gcc ccc  
 aat ccc tgt ata gaa tgt ggc caa aag tcc ctc cac ctg 931 His Ala Pro Asn Pro Cys Ile Glu Cys Gly Gln  
 Lys Ser Leu His Leu260 265 270 275cac cct gag acc ttc cat cacctgagc cat ctg gaa ggc ctg gtg ctg  
 979 His Pro Glu Thr Phe His His Leu Ser His Leu Glu Gly Leu Val Leu 280 285 290aag gac agc tct  
 ctc cat aca ctg aac tct tcc tgg ttc caa ggt ctg 1027 Lys Asp Ser Ser Leu His Thr Leu Asn Ser Ser  
 Trp Phe Gln Gly Leu 295 300 305 gtc aac ctc tcg gtgctg gac cta agc gag aac ttt ctc tat gaa agc 1075  
 Val Asn Leu Ser Val Leu Asp Leu Ser Glu Asn Phe Leu Tyr Glu Ser 310 315 320 atc aac cac acc aat  
 gccttt cag aac cta acc cgc ctg cgc aag ctc 1123 Ile Asn His ThrAsn Ala Phe Gln Asn Leu Thr Arg  
 Leu Arg Lys Leu 325 330 335 aac ctg tcc ttc aat tac cgcaag aag gta tcc ttt gcc cgc ctc cac 1171  
 Asn Leu Ser Phe Asn Tyr Arg Lys Lys Val Ser Phe Ala Arg Leu His340 345 350 355ctg gca agt tcc  
 ttc aag aac ctg gtg tca ctg cag gag ctg aac atg 1219 Leu Ala Ser Ser Phe Lys Asn Leu Val Ser Leu  
 Gln Glu Leu Asn Met 360 365 370 aac ggc atc ttc ttc cgc tcg ctc aac aag tac acg ctc aga tgg ctg  
 1267 Asn Gly Ile Phe Phe Arg Ser Leu Asn Lys Tyr Thr Leu Arg Trp Leu 375 380 385 gcc gat ctg  
 ccc aaactc cac act ctg cat ctt caa atg aacttc atc 1315 Ala Asp Leu Pro Lys Leu His Thr Leu His  
 Leu Gln Met Asn Phe Ile 390 395 400 aac cag gca cag ctc agcatc ttt ggt acc ttc cga gcc ctt cgc ttt  
 1363 A sn Gln Ala Gln Leu Ser-Ile-Phe-Gly-Thr Phe Arg Ala Leu Arg Phe 405 410 415 gtg gac ttg  
 tca gac-aat-cgc-atc-agt ggg cct tca acg ctg tca gaa 1411 Val Asp Leu Ser Asp Asn Arg Ile Ser Gly  
 Pro Ser Thr Leu Ser Glu420 425 430 435gcc acc cct gaa gag gca gat gat gca gag caggaggag ctg ttg  
 tct 1459 Ala Thr Pro Glu Glu Ala Asp Asp Ala Glu Gln Glu Glu Leu Leu Ser 440 445 450 gcg gat cct  
 cac cca gct cca ctg agc acc cct gct tct aag aac ttc 1507 Ala Asp Pro His Pro Ala Pro Leu Ser Thr

Pro Ala Ser Lys Asn Phe 455 460 465 atg gac agg tgt aag aac ttc aag ttc acc atg gac ctg tct cgg aac  
 1555 Met Asp Arg Cys Lys Asn Phe Lys Phe Thr Met Asp Leu Ser Arg Asn 470 475 480 aac ctg gtg  
 act atc aag cca gag atg ttt gtc aat ctc tca cgc ctc 1603 Asn Leu Val Thr Ile Lys Pro Glu Met Phe  
 Val Asn Leu Ser Arg Leu 485 490 495 cag tgt ctt agc ctg agc cac aac tcc att gca cag gct gtc aat ggc  
 1651 Gln Cys Leu Ser Leu Ser His Asn Ser Ile Ala Gln Ala Val Asn Gly 500 505 510 515 tct cag ttc ctg  
 ccg ctg actaatctg cag gtg ctg gac ctg tcc cat 1699 Ser Gln Phe Leu Pro Leu Thr Asn Leu Gln Val  
 Leu Asp Leu Ser His 520 525 530 aac aaa ctg gac ttg tac cac tgg aaa tgg ttc agt gag cta cca cag  
 1747 Asn Lys Leu Asp Leu Tyr His Trp Lys Ser Phe Ser Glu Leu Pro Gln 535 540 545 ttg cag gcc  
 ctg gac ctg agc tac aac agc cag ccc ttt agc atg aag 1795 Leu Gln Ala Leu Asp Leu Ser Tyr Asn Ser  
 Gln Pro Phe Ser Met Lys 550 555 560 ggt ata ggc cac aat ttc agt ttt gtg gcc cat ctg tcc atg cta cac  
 1843 Gly Ile Gly His Asn Phe Ser Phe Val Ala His Leu Ser Met Leu His 565 570 575 agc ctt agc ctg  
 gca cac aat gac att cat acc cgt gtg tcc tca cat 1891 Ser Leu Ser Leu Ala His Asn Asp Ile His Thr  
 Arg Val Ser Ser His 580 585 590 595 ctc aac agc aac tca gtg aggtttctt gac ttc agc ggc aac ggt atg  
 1939 Leu Asn Ser Asn Ser Val Arg Phe Leu Asp Phe Ser Gly Asn Gly Met 600 605 610 ggc cgc atg  
 tgg gat gag ggg gcc ctt tat ctc cat ttc ttc caa gcc 1987 Gly-Arg-Met-Trp-Asp Glu Gly Gly Leu  
 Tyr-Leu-His-Phe-Phe-Gln-Gly 615 620 625 ctg agt ggc ctg ctg aag ctg gac ctg tct caa aat aac ctg  
 cat atc 2035 Leu Ser Gly Leu Leu Lys Leu Asp Leu Ser Gln Asn Asn Leu His 630 635 640  
 ctc cgg ccc cag aac ctt gac aac ctc ccc aag agc ctg aag ctg ctg 2083 Leu Arg Pro Gln Asn Leu Asp  
 Asn Leu Pro Lys Ser Leu Lys Leu Leu 645 650 655 agc ctc cga gac aac tac cta tct ttc ttt aac tgg  
 acc agt ctg tcc 2131 Ser Leu Arg Asp Asn Tyr Leu Ser Phe Phe Asn Trp Thr Ser Leu Ser 660 665  
 670 675 ttc ctg ccc aac ctg gaa gtctagac ctg gca gcc aac cag cta aag 2179 Phe Leu Pro Asn Leu  
 Glu Val Leu Asp Leu Ala Gly Asn Gln Leu Lys 680 685 690 gcc ctg acc aat ggc acc ctg cct aat ggc  
 acc ctc ctc cag aaa ctg 2227 Ala Leu Thr Asn Gly Thr Leu Pro Asn Gly Thr Leu Leu Gln Lys Leu  
 695 700 705 gat gtc agc agc aac agt atc gtc tct gtg gtc cca gcc ttc ttc gct 2275 Asp Val Ser Ser  
 Asn Ser Ile Val Ser Val Val Pro Ala Phe Phe Ala 710 715 720 ctg gcg gtc gag ctg aaa gag gtc aac ctc  
 agc cac aac att ctc aag 2323 Leu Ala Val Glu Leu Lys Glu Val Asn Leu Ser His Asn Ile Leu Lys 725  
 730 735 acg gtg gat cgc tcc tgg ttt ggg ccc att gtg atg aac ctg aca gtt 2371 Thr Val Asp Arg Ser Trp  
 Phe Gly Pro Ile Val Met Asn Leu Thr Val 740 745 750 755 cta gac gtg aga agc aac cctctgcac tgt gcc  
 tgt ggg gca gcc ttc 2419 Leu Asp Val Arg Ser Asn Pro Leu His Cys Ala Cys Gly Ala Ala Phe 760 765  
 770 gta gac tta ctg ttg gag gtg cag acc aag gtg cct gcc ctg gct aat 2467 Val Asp Leu Leu Leu Glu Val  
 Gln Thr Lys Val Pro Gly Leu Ala Asn 775 780 785 ggt gtg aag tgt ggc agc ccc gcc cag ctg cag gcc  
 cgt agc atc ttc 2515 Gly Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Arg Ser Ile Phe 790 795 800  
 gca cag gac ctg cgg ctg tgc ctg gat gag gtc ctc tct tgg gac tgc 2563 Ala Gln Asp Leu Arg Leu Cys  
 Leu Asp Glu Val Leu Ser Trp Asp Cys 805 810 815 ttt ggc ctt tca ctc ttg gct gtg gcc gtg ggc atg gtg  
 gtg cct ata 2611 Phe Gly Leu Ser Leu Leu Ala Val Ala-Val-Gly-Met-Val Val Pro Ile 820 825 830  
 835 ctg cac cat ctc tgc gcc tgg gac gtc tgg tac tgt ttt cat ctg tgc 2659 Leu His His Leu Cys Gly Trp  
 Asp Val Trp Tyr Cys Phe His Leu Cys 840 845 850 ctg gca tgg cta cct ttg ctg gcc cgc agc cga cgc  
 agc gcc caa gct 2707 Leu Ala Trp Leu Pro Leu Leu Ala Arg Ser Arg Arg Ser Ala Gln Ala 855 860 865  
 ctc ccc tat gat gccttc gtg gtg ttc gat aag gca cag agcgca gtt 2755 Leu Pro Tyr Asp Ala Phe Val Val  
 Phe Asp Lys Ala Gln Ser Ala Val 870 875 880 gcg gac tgg gtg tat aacgag ctg cgg gtg cgg ctg gag gag  
 cgg cgc 2803 Ala Asp Trp Val Tyr Asn Glu Leu Arg Val Arg Leu Glu Glu Arg Arg 885 890 895 ggt cgc  
 cga gcccta cgc ttgtgt ctg gag gac cga gat tgg ctg cct 2851 Gly Arg Arg Ala Leu Arg Leu Cys Leu Glu  
 Asp Arg Asp Trp Leu Pro 900 905 910 915 ggc cag acg ctc ttc gag aac ctc tgg gct tcc atc tat ggg agc  
 cgc 2899 Gly Gln Thr Leu Phe Glu Asn Leu Trp Ala Ser Ile Tyr Gly Ser Arg 920 925 930 aag act cta  
 ttt gtg ctg gcc cac acg gac cgc gtc agt ggc ctc ctg 2947 Lys Thr Leu Phe Val Leu Ala His Thr Asp  
 Arg Val Ser Gly Leu Leu 935 940 945 cgc acc agc ttc ctgctg gct cag cag cgc ctg ttg gaa gaccgc aag  
 2995 Arg Thr Ser Phe Leu Leu Ala Gln Gln Arg Leu Leu Glu Asp Arg Lys 950 955 960 gac gtg gtg gtg  
 ttg gtgac ctg cgt ccg gat gcc cac cgc tcc cgc 3043 Asp Val Val Val Leu Val Ile Leu Arg Pro Asp Ala  
 His Arg Ser Arg 965 970 975 tat gtg cga ctgcgc cag cgtctc tgc cgc cag agt gtg ctc ttc tgg 3091 Tyr  
 Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val Leu Phe Trp 980 985 990 995 ccc cag cag ccc aac  
 ggg cag ggg gcc ttc tgg gcc cag ctg agt aca 3139 Pro Gln Gln Pro Asn Gly Gln Gly Gly Phe Trp Ala  
 Gln Leu Ser Thr 1000 1005 1010 gcc ctg act agg gac-aac-cgc-cac-ttc tat aac cag aac ttc tgc cgg  
 3187 Ala Leu Thr Arg Asp Asn-Arg-His-Phe-Tyr Asn Gln Asn Phe Cys-Arg 1015 1020 1025 gga cct  
 aca gca-gaa-tag-ctcagagcaa cagctggaaa cagctgcatc 3235 Gly Pro Thr Ala Glu 1030 ttcctgcctg



gttcccgagt tgcctctgcct gcccttgcctct gtcttactac accgctattt 3295 ggcaagtgcg caatatatgc  
 taccaagccaccaggcccccac ggagcaaagg ttggcagtaa 3355 aggtagttt tcttcccatg catctttcag gagagtgaag  
 atagacacca gacccacaca 3415 gaacaggact ggagttcattctctgcccct ccacccact ttgcctgtct ctgtat 3471  
 <210> 4 <211> 1032<212> PRT <213> Mus musculus <400> 4 Met Val Leu Arg Arg Arg ThrLeu His  
 Pro Leu Ser Leu Leu Val Gln 1 5 10 15 Ala Ala Val Leu Ala Glu Thr Leu Ala Leu Gly Thr Leu Pro Ala  
 Phe 20 25 30 Leu Pro Cys Glu Leu Lys Pro His Gly Leu Val Asp Cys Asn Trp Leu 35 40 45 Phe Leu  
 Lys Ser Val Pro ArgPhe Ser Ala Ala Ala Ser Cys Ser Asn 50 55 60 Ile ThrArg LeuSer Leu Ile Ser Asn  
 Arg Ile His His Leu His Asn 65 70 75 80 Ser Asp Phe Val His Leu Ser Asn Leu Arg Gln Leu Asn Leu  
 Lys Trp 85 90 95 Asn Cys Pro Pro Thr Gly Leu Ser Pro Leu His Phe Ser Cys His Met 100 105 110  
 Thr Ile Glu Pro Arg Thr Phe LeuAla Met Arg Thr Leu Glu Glu Leu 115 120 125 Asn Leu Ser Tyr Asn  
 GlyIle Thr Thr Val Pro Arg Leu Pro Ser Ser 130 135 140 Leu Val Asn Leu Ser Leu Ser His Thr AsnIle  
 Leu Val Leu Asp Ala 145 150 155 160 Asn Ser Leu Ala Gly Leu Tyr Ser Leu Arg Val Leu Phe Met  
 Asp Gly 165 170 175 Asn Cys Tyr Tyr Lys Asn Pro Cys Thr Gly Ala Val Lys Val Thr Pro 180 185 190  
 Gly Ala Leu Leu Gly Leu Ser Asn Leu Thr His Leu Ser Val Lys Tyr 195 200 205 Asn Asn Leu Thr Lys  
 Val Pro Arg Gln Leu Pro Pro Ser Leu Glu Tyr 210 215 220 Leu Leu Val Ser Tyr Asn Leu Ile  
 Val-Lys-Leu-Gly-Pro-Glu Asp Leu 225 230 235 240 Ala Asn Leu Thr Ser Leu Arg Val  
 Leu-Asp-Val-Gly-Gly-Asn-Cys-Arg 245 250 255 Arg Cys Asp His Ala Pro Asn Pro Cys Ile Glu Cys  
 Gly Gln Lys Ser 260 265270 Leu His Leu His Pro Glu Thr Phe His His Leu Ser His Leu Glu Gly 275  
 280 285 Leu Val Leu Lys Asp Ser Ser Leu His Thr Leu Asn Ser Ser Trp Phe 290 295 300 Gln Gly  
 Leu Val Asn Leu Ser Val Leu Asp Leu Ser Glu Asn Phe Leu 305 310 315 320 Tyr Glu Ser Ile Asn His  
 Thr Asn Ala Phe Gln Asn Leu Thr Arg Leu 325 330 335 Arg Lys Leu Asn Leu Ser Phe Asn Tyr Arg  
 Lys Lys Val Ser Phe Ala 340 345 350 Arg Leu His Leu Ala Ser Ser Phe Lys Asn Leu Val Ser Leu Gln  
 Glu 355 360 365 Leu Asn Met Asn Gly Ile Phe Phe Arg Ser Leu Asn Lys Tyr Thr Leu 370 375 380  
 Arg Trp Leu Ala Asp Leu Pro Lys Leu His Thr Leu His Leu Gln Met 385 390 395 400 Asn Phe Ile Asn  
 Gln Ala Gln Leu Ser Ile Phe Gly Thr Phe Arg Ala 405 410 415 Leu Arg Phe Val Asp Leu Ser Asp Asn  
 Arg Ile Ser Gly Pro Ser Thr 420 425 430 Leu Ser Glu Ala Thr Pro Glu Glu Ala Asp Asp Ala Glu Gln  
 Glu Glu 435 440 445 Leu Leu Ser Ala Asp Pro His Pro Ala Pro Leu Ser Thr Pro Ala Ser 450 455 460  
 Lys Asn Phe Met Asp Arg Cys Lys Asn Phe Lys Phe Thr Met Asp Leu 465 470 475 480 Ser Arg Asn  
 Asn Leu Val Thr Ile Lys Pro Glu Met Phe Val Asn Leu 485 490 495 Ser Arg Leu Gln Cys Leu Ser Leu  
 Ser His Asn Ser Ile Ala Gln Ala 500 505 510 Val Asn Gly Ser Gln Phe Leu Pro Leu Thr Asn Leu Gln  
 Val Leu Asp 515 520 525 Leu Ser His Asn Lys Leu Asp Leu Tyr His Trp Lys Ser Phe Ser Glu 530  
 535 540 Leu Pro Gln Leu Gln Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe 545 550 555 560 Ser  
 Met Lys Gly Ile Gly His Asn Phe-Ser-Phe-Val-Ala-His-Leu-Ser 565 570 575 Met Leu His Ser Leu  
 Ser Leu Ala His-Asn-Asp-Ile-His-Thr-Arg-Val 580 585 590 Ser Ser His Leu Asn Ser Asn Ser Val  
 Arg Phe Leu Asp Phe Ser Gly 595 600 605 Asn Gly Met Gly Arg Met Trp Asp Glu Gly Gly Leu Tyr  
 Leu His Phe 610 615 620 Phe Gln Gly Leu Ser Gly Leu Leu Lys Leu Asp Leu Ser Gln Asn Asn 625  
 630 635 640 Leu His Ile Leu Arg Pro Gln Asn Leu Asp Asn Leu Pro Lys Ser Leu 645 650 655 Lys  
 Leu Leu Ser Leu Arg Asp Asn Tyr Leu Ser Phe Phe Asn Trp Thr 660 665 670 Ser Leu Ser Phe Leu  
 Pro Asn Leu Glu Val Leu Asp Leu Ala Gly Asn 675 680 685 Gln Leu Lys Ala Leu Thr Asn Gly Thr Leu  
 Pro Asn Gly Thr Leu Leu 690 695 700 Gln Lys Leu Asp Val Ser Ser Asn Ser Ile Val Ser Val Val Pro  
 Ala 705 710 715 720 Phe Phe Ala Leu Ala Val Glu Leu Lys Glu Val Asn Leu Ser His Asn 725 730 735  
 Ile Leu Lys Thr Val Asp Arg Ser Trp Phe Gly Pro Ile Val Met Asn 740 745 750 Leu Thr Val Leu Asp  
 Val Arg Ser Asn Pro Leu His Cys Ala Cys Gly 755 760 765 Ala Ala Phe Val Asp Leu Leu Leu Glu Val  
 Gln Thr Lys Val Pro Gly 770 775 780 Leu Ala Asn Gly Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly  
 Arg 785 790 795 800 Ser Ile Phe Ala Gln Asp Leu Arg Leu Cys Leu Asp Glu Val Leu Ser 805 810 815  
 Trp Asp Cys Phe Gly Leu Ser Leu Leu Ala Val Ala Val Gly Met Val 820 825 830 Val Pro Ile Leu His  
 His Leu Cys Gly Trp Asp Val Trp Tyr Cys Phe 835 840 845 His Leu Cys Leu Ala Trp Leu Pro Leu  
 Leu Ala Arg Ser Arg Arg Ser 850 855 860 Ala Gln Ala Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys  
 Ala Gln 865 870 875 880 Ser Ala Val Ala Asp Trp Val Tyr Asn Glu Leu Arg Val Arg Leu Glu 885 890  
 895 Glu Arg Arg Gly Arg Arg Ala Leu Arg-Leu-Cys-Leu-Glu-Asp-Arg-Asp 900 905 910 Trp Leu Pro  
 Gly Gln Thr Leu Phe GluAsn-Leu-Trp-Ala-Ser-Ile-Tyr 915 920925 Gly Ser Arg Lys Thr Leu Phe Val  
 Leu Ala His Thr Asp Arg Val Ser 930 935 940 Gly Leu Leu Arg Thr Ser Phe Leu Leu Ala Gln Gln Arg  
 Leu Leu Glu 945 950 955 960 Asp Arg Lys Asp ValVal Val Leu Val Ile Leu Arg Pro Asp Ala His 965  
 970 975 Arg Ser Arg Tyr Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val 980 985 990 Leu Phe Trp

41  
Pro Gln Gln Pro Asn Gly Gln Gly Gly Phe Trp Ala Gln 995 1000 1005 Leu Ser Thr AlaLeu ThrArg Asp  
Asn Arg His Phe Tyr Asn Gln Asn 1010 1015 1020 Phe Cys ArgGly Pro Thr Ala Glu 1025 1030 <210>  
5 <211> 20<212> DNA <213> Artificial Sequence<220> <223> Description of Artificial Sequence:CpG  
ODN <400> 5 tccatgacgt tcctgatgct 20

---

[Translation done.]

\* NOTICES \*

JPO and NCIPi are not responsible for any damages caused by the use of this translation.

- 1.This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.\*\*\*\* shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

---

DESCRIPTION OF DRAWINGS

---

[Brief Description of the Drawings]

[Drawing 1] It is drawing showing the genetic map of TLR9 knockout mouse of this invention, and a wild type mouse.

[Drawing 2] It is drawing showing the result of the Southern blot analysis of TLR9 knockout mouse of this invention.

[Drawing 3] It is drawing showing the result of the Northern blot analysis in the spleen cell of TLR9 knockout mouse of this invention.

[Drawing 4] It is drawing showing the comparison result of the amino acid sequence of TLR9 knockout mouse of this invention, and a wild type mouse.

[Drawing 5] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of TNFalpha by ODN, PGN, or LPS induction, IL-6, or the amount of production of IL12.

[Drawing 6] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of the cell proliferation response by ODN or LPS induction.

[Drawing 7] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of the amount of production of IL-12 by ODN or LPS induction.

[Drawing 8] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of the amount of manifestations of the CD40, CD80, CD86, and the MHC class II by ODN or LPS induction.

[Drawing 9] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of activation of NF-kappa B by ODN or LPS induction.

[Drawing 10] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of the activation of JNK by ODN or LPS induction.

[Drawing 11] CpG in TLR9 knockout mouse and wild type mouse of this invention It is drawing showing the result of activation of ODN or IRAK by LPS induction.

---

[Translation done.]

(19) 日本国特許庁 (J P)

(12) 公開特許公報 (A)

(11) 特許出願公開番号

特開2002-34565

(P 2 0 0 2 - 3 4 5 6 5 A)

(43) 公開日 平成14年2月5日(2002.2.5)

(51) Int. Cl. <sup>7</sup>	識別記号	F I	ターマコード (参考)
C12N 15/09	ZNA	A61K 45/00	2G045
A61K 38/00		A61P 31/04	4B024
45/00		35/00	4B063
A61P 31/04		37/08	4B064
35/00		C07K 14/005	4B065

審査請求 未請求 請求項の数30 O L (全32頁) 最終頁に続く

(21) 出願番号	特願2000-219652 (P 2000-219652)	(71) 出願人	396020800 科学技術振興事業団 埼玉県川口市本町4丁目1番8号
(22) 出願日	平成12年7月19日(2000.7.19)	(72) 発明者	審良 静男 大阪府高槻市辻子1丁目7-16
		(72) 発明者	辺見 弘明 大阪府茨木市北春日丘4-11-47-112
		(74) 代理人	100107984 弁理士 廣田 雅紀

最終頁に続く

(54) 【発明の名称】 細菌DNAを特異的に認識する受容体タンパク質

(57) 【要約】

【課題】 非メチル化C p G配列を有する細菌DNAを特異的に認識する受容体タンパク質や、それをコードする遺伝子DNAや、細菌性伝染病に対する宿主免疫細胞の応答性を調べる上で有用な実験モデル動物を提供すること。

【解決手段】 非メチル化C p G配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAを、BLASTサーチによりスクリーニングし、各種TLRと高い相似性を有する多くのESTクローンをスクリーニングし、これらをプローブにして、マウス・マクロファージcDNAライブラリーから完全長cDNAを単離し、cDNAの塩基配列を解析してLRR及びTIR領域などの保存領域が存在するTLR9であることを確認した後、ノックアウトマウスを作製し、TLR9が細菌DNAの非メチル化C p G配列を含むオリゴヌクレオチドの受容体タンパク質であることを確認した。

## 【特許請求の範囲】

【請求項 1】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA。

【請求項 2】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質が、以下の (a) 又は (b) のタンパク質であることを特徴とする請求項 1 記載の DNA。

(a) 配列番号 2 に示されるアミノ酸配列からなるタンパク質

(b) 配列番号 2 に示されるアミノ酸配列において、1 若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ非メチル化 C p G 配列を有する細菌 DNA に対して反応性を有するタンパク質

【請求項 3】 配列番号 1 に示される塩基配列又はその相補的配列並びにこれらの配列の一部または全部を含むことを特徴とする請求項 1 記載の DNA。

【請求項 4】 請求項 3 記載の遺伝子を構成する DNA とストリンジェントな条件下でハイブリダイズすることを特徴とする請求項 1 記載の DNA。

【請求項 5】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質が、以下の (a) 又は (b) のタンパク質であることを特徴とする請求項 1 記載の DNA。

(a) 配列番号 4 に示されるアミノ酸配列からなるタンパク質

(b) 配列番号 4 に示されるアミノ酸配列において、1 若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ非メチル化 C p G 配列を有する細菌 DNA に対して反応性を有するタンパク質

【請求項 6】 配列番号 3 に示される塩基配列又はその相補的配列並びにこれらの配列の一部または全部を含むことを特徴とする請求項 1 記載の DNA。

【請求項 7】 請求項 6 記載の遺伝子を構成する DNA とストリンジェントな条件下でハイブリダイズすることを特徴とする請求項 1 記載の DNA。

【請求項 8】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質。

【請求項 9】 配列番号 2 に示されるアミノ酸配列からなることを特徴とする請求項 8 記載のタンパク質。

【請求項 10】 配列番号 2 に示されるアミノ酸配列において、1 若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなることを特徴とする請求項 8 記載のタンパク質。

【請求項 11】 配列番号 4 に示されるアミノ酸配列からなることを特徴とする請求項 8 記載のタンパク質。

【請求項 12】 配列番号 4 に示されるアミノ酸配列において、1 若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなることを特徴とする請求項 8 記載のタンパク質。

【請求項 13】 請求項 8～12 のいずれか記載のタンパク質と、マーカータンパク質及び／又はペプチドタグとを結合させた融合タンパク質。

【請求項 14】 請求項 8～12 のいずれか記載のタンパク質と特異的に結合する抗体。

【請求項 15】 抗体がモノクローナル抗体であることを特徴とする請求項 14 記載の抗体。

【請求項 16】 請求項 8～12 のいずれか記載のタンパク質を発現することができる発現系を含んでなる宿主細胞。

【請求項 17】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現することを特徴とする非ヒト動物。

【請求項 18】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損したことを特徴とする非ヒト動物。

【請求項 19】 非メチル化 C p G 配列を有する細菌 DNA に対して不反応性であることを特徴とする請求項 18 記載の非ヒト動物。

【請求項 20】 齧歯目動物が、マウスであることを特徴とする請求項 17～19 のいずれか記載の非ヒト動物。

【請求項 21】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した細胞に、請求項 1～7 のいずれか記載の DNA を導入することを特徴とする非メチル化 C p G 配列を有する細菌 DNA に対して反応性を有するタンパク質を発現する細胞の調製方法。

【請求項 22】 請求項 21 記載の非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を発現する細胞の調製方法により得られることを特徴とする非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を発現する細胞。

【請求項 23】 被検物質の存在下、非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を発現している細胞をインビトロで培養し、TLR9 活性を測定・評価することを特徴とする非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法。

【請求項 24】 非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物に被検物質を投与し、該非ヒト動物から得られるマクロファージ又は脾臓細胞の TLR9 活性を測定・評価することを特徴とする非メチル化 C p G 配列を有する細菌 DNA を特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法。

【請求項 25】 非メチル化 C p G 配列を有する細菌 D

10

20

30

40

50

NAを特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現した非ヒト動物に被検物質を投与し、該非ヒト動物から得られるマクロファージ又は脾臓細胞のTLR9活性を測定・評価することを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法。

【請求項26】 非ヒト動物が、マウスであることを特徴とする請求項24又は25記載の非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質のアゴニスト又はアンタゴニストのスクリーニング方法。

【請求項27】 請求項23～26のいずれか記載の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法により得られる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニスト。

【請求項28】 非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の全部又はその一部を有効成分として含有する医薬組成物。

【請求項29】 請求項27記載のアゴニスト又はアンタゴニストを有効成分として含有する医薬組成物。

【請求項30】 検体中の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAと、請求項3記載のDNAとの塩基配列を比較することができる、請求項3記載のDNAを含むことを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA配列の欠失、置換及び／又は付加に関連する疾病の診断キット。

【発明の詳細な説明】

【0001】

【発明の属する技術分野】本発明は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質、該受容体タンパク質の遺伝子及びそれらの利用に関する。

【0002】

【従来の技術】ツール(Toll)遺伝子は、ショウジョウバエの胚発生中の背腹軸の決定(Cell 52, 269-279, 1988, Annu. Rev. Cell Dev. Biol. 12, 393-416, 1996)、また成体における抗真菌性免疫応答に必要であることが知られている(Cell 86, 973-983, 1996)。かかるTollは、細胞外領域にロイシンリッチリピート(LRR)を有するI型膜貫通受容体であり、この細胞質内領域は、哺乳類インターロイキン-1受容体(IL-1R)の細胞質内領域と相同性が高いことが明らかとなっている(Nature 351, 355-356, 1991, Annu. Rev. Cell Dev. Biol. 12, 393-416, 1996, J. Leukoc. Biol. 63, 650-657, 1998)。

【0003】近年、Toll様受容体(TLR)と呼ばれるTollの哺乳類のホモログが同定され、TLR2やTLR4など現在までに6つのファミリーが報告されている(Nature 388, 394-397, 1997, Proc. Natl. Acad. Sci. USA 95, 588-593, 1998, Blood 91, 4020-4027, 1998, Gene 231, 59-65, 1999)。このTLRファミリーは、上記IL-1Rと同様にアダプタータンパク質であるMyD88を介し、IL-1R結合キナーゼ(IRAK)をリクルートし、TRAF6を活性化し、下流のNF- $\kappa$ Bを活性化することが知られている(J. Exp. Med. 187, 2097-2101, 1998, Mol. Cell 2, 253-258, 1998, Immunity 11, 115-122, 1999)。また、哺乳類におけるTLRファミリーの役割は、細菌の共通構造を認識するパターン認識受容体(PRR: pattern recognition receptor)として、先天的な免疫認識に関わっているとも考えられている(Cell 91, 295-298, 1997)。

【0004】上記PRRにより認識される病原体会合分子パターン(PAMP: pathogen-associated molecular pattern)の一つは、グラム陰性菌の外膜の主成分であるリポ多糖(LPS)であって(Cell 91, 295-298, 1997)、かかるLPSが宿主細胞を刺激して宿主細胞にTNF $\alpha$ 、IL-1及びIL-6等の各種炎症性サイトカインを産生させること(Adv. Immunol. 28, 293-450, 1979, Annu. Rev. Immunol. 13, 437-457, 1995)や、LPS結合タンパク質(LBP: LPS-binding protein)により捕獲されたLPSが細胞表面上のCD14に引き渡されることが知られている(Science 249, 1431-1433, 1990, Annu. Rev. Immunol. 13, 437-457, 1995)。本発明者らは、TLR4のノックアウトマウスを作製し、TLR4ノックアウトウスが上記グラム陰性菌の外膜の主成分であるLPSに不応答性であること(J. Immunol. 162, 3749-3752, 1999)や、TLR2ノックアウトマウスを作製し、TLR2ノックアウトマウスのマクロファージがグラム陽性菌細胞壁やその構成成分であるペプチドグリカンに対する反応性が低下すること(Immunity, 11, 443-451, 1999)を報告している。

【0005】他方、細菌DNA(バクテリア由来DNA)や非メチル化CpG配列を含むオリゴヌクレオチドが、マウス及びヒトの免疫細胞を刺激すること(Trends Microbiol. 4, 73-76, 1996, Trends Microbiol. 6, 496-500, 1998)や、IL-12及びIFN $\gamma$ の放出に支配されるTヘルパー1細胞(Th1)様炎症性応答を刺激すること(EMBO J. 18, 6973-6982, 1999, J. Immunol. 161, 3042-3049, 1998, Proc. Natl. Acad. Sci. USA 96, 9305-9310, 1999)から、CpG配列を含むオリゴヌクレオチドは、癌、アレルギー及び伝染病のワクチンを含むワクチン戦略のアジュバントとしての使用可能性が提唱されている(Adv. Immunol. 73, 329-368, 1999, Curr. Opin. Immunol. 12, 35-43, 2000, Immunity 11, 1

23-129, 1999). このように臨床実用において効果が期待されるにも関わらず、非メチル化CpG配列を含む細菌DNAが免疫細胞を活性化する分子メカニズムはよくわかっていない。

#### 【0006】

【発明が解決しようとする課題】上記のように、メチル化されていないCpGモチーフを含有するバクテリア由来DNAは免疫細胞を非常に活性化し、Th1の応答を誘導するが、その分子レベルでの活動はあまり理解されていない。本発明の課題は、細菌DNAの非メチル化CpG配列を含むオリゴヌクレオチドの分子レベルでの作用を明らかにすることができる、非メチル化CpG配列を有する細菌DNAを特異的に認識するTLRファミリーのメンバー受容体タンパク質TLR9や、それをコードするDNAや、細菌性伝染病に対する宿主免疫細胞の応答性を調べる上で有用な実験モデル動物を提供することにある。

#### 【0007】

【課題を解決するための手段】細菌の共通構造を認識するパターン認識受容体として、先天的な免疫認識に関わっている哺乳類におけるTLRファミリーは、現在までに6つのメンバー(TLR1-6)が公表されており(Nature 388, 394-397, 1997, Proc. Natl. Acad. Sci. USA, 95, 588-593, 1998, Gene 231, 59-65, 1999)、TLR7及びTLR8の新たな2つのメンバーがGenBankに登録されている(登録番号AF240467及びAF246971)。また、TLR9についても完全長cDNAが見い出されGenBankに登録されている(登録番号AF245704)が、その機能については知られていなかった。

【0008】本発明者らは、非メチル化CpG配列を有する細菌DNAを特異的に認識するTLRファミリーのメンバー受容体タンパク質をコードするDNAを、BLASTサーチによりスクリーニングし、既に同定されている各種TLRと高い相似性を有する多くのシーケン・タグ(EST)クローンをスクリーニングし、これらの遺伝子フラグメントをプローブにして、マウス・マクロファージcDNAライブラリーから完全な長さを有するcDNAを単離し、これを用いてヒトcDNAも単離した。次に、これらcDNAの塩基配列を解析し、このTLRファミリーにLRP及びTIR領域などの保存領域が存在するTLR9であることを確認した。そこで、このTLR9ノックアウトマウスを作製し、TLR9が細菌DNAの非メチル化CpG配列を含むオリゴヌクレオチドの受容体タンパク質であることを明らかにし、本発明を完成するに至った。

【0009】すなわち本発明は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA(請求項1)や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タ

ンパク質が、以下の(a)又は(b)のタンパク質であることを特徴とする請求項1記載のDNA(a)配列番号2に示されるアミノ酸配列からなるタンパク質(b)配列番号2に示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質(請求項2)や、配列番号1に示される塩基配列又はその相補的配列並びにこれらの配列の一部または全部を含むことを特徴とする請求項1記載のDNA(請求項3)や、請求項3記載の遺伝子を構成するDNAとストリンジントな条件下でハイブリダイズすることを特徴とする請求項1記載のDNA(請求項4)や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質が、以下の(a)又は(b)のタンパク質であることを特徴とする請求項1記載のDNA(a)配列番号4に示されるアミノ酸配列からなるタンパク質(b)配列番号4に示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質(請求項5)や、配列番号3に示される塩基配列又はその相補的配列並びにこれらの配列の一部または全部を含むことを特徴とする請求項1記載のDNA(請求項6)や、請求項6記載の遺伝子を構成するDNAとストリンジントな条件下でハイブリダイズすることを特徴とする請求項1記載のDNA(請求項7)に関する。

【0010】また本発明は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質(請求項8)や、配列番号2に示されるアミノ酸配列からなることを特徴とする請求項8記載のタンパク質(請求項9)や、配列番号2に示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなることを特徴とする請求項8記載のタンパク質(請求項10)や、配列番号4に示されるアミノ酸配列からなることを特徴とする請求項8記載のタンパク質(請求項11)や、配列番号4に示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなることを特徴とする請求項8記載のタンパク質(請求項12)に関する。

【0011】また本発明は、請求項8～12のいずれか記載のタンパク質と、マーカータンパク質及び／又はペプチドタグとを結合させた融合タンパク質(請求項13)や、請求項8～12のいずれか記載のタンパク質と特異的に結合する抗体(請求項14)や、抗体がモノクローナル抗体であることを特徴とする請求項14記載の抗体(請求項15)や、請求項8～12のいずれか記載のタンパク質を発現することができる発現系を含んでなる宿主細胞(請求項16)に関する。

【0012】また本発明は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現することを特徴とする非ヒト動物（請求項17）や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損したことを特徴とする非ヒト動物（請求項18）や、非メチル化CpG配列を有する細菌DNAに対して不反応性であることを特徴とする請求項18記載の非ヒト動物（請求項19）や、齧歯目動物が、マウスであることを特徴とする請求項17～19のいずれか記載の非ヒト動物（請求項20）に関する。

【0013】また本発明は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した細胞に、請求項1～7のいずれか記載のDNAを導入することを特徴とする非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質を発現する細胞の調製方法（請求項21）や、請求項21記載の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞の調製方法により得られることを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞（請求項22）に関する。

【0014】また本発明は、被検物質の存在下、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現している細胞をインビトロで培養し、TLR9活性を測定・評価することを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法（請求項23）や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物に被検物質を投与し、該非ヒト動物から得られるマクロファージ又は脾臓細胞のTLR9活性を測定・評価することを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法（請求項24）や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現した非ヒト動物に被検物質を投与し、該非ヒト動物から得られるマクロファージ又は脾臓細胞のTLR9活性を測定・評価することを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法（請求項25）や、非ヒト動物が、マウスであることを特徴とする請求項24又は25記載の非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質のアゴニスト又はアンタゴニストのスクリーニング方法（請求項26）に関する。

【0015】また本発明は、請求項23～26のいずれか記載の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法により得られる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニスト（請求項27）や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の全部又はその一部を有効成分として含有する医薬組成物（請求項28）や、請求項27記載のアゴニスト又はアンタゴニストを有効成分として含有する医薬組成物（請求項29）や、検体中の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAと、請求項3記載のDNAとの塩基配列を比較することができる、請求項3記載のDNAを含むことを特徴とする非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA配列の欠失、置換及び／又は付加に関連する疾病の診断キット（請求項30）に関する。

#### 【0016】

【発明の実施の形態】本発明における非メチル化CpG配列を有する細菌DNAとしては、T細胞、B細胞、抗原提示細胞等の免疫細胞を活性化し、免疫応答を誘導することができる、メチル化されていないCpGモチーフを有するオリゴデオキシヌクレオチド（ODN）等のバクテリアに由来するDNAであればどのようなものでもよく、エセリシア・コリ、クレブシエラ・ニューモニエ、シュドモナス・アエルギノサ、サルモネラ・チフィウム、セラチア・マルセッセンス、フレクサナー赤痢菌、ビブリオ・コレレエ、サルモネラ・ミネソタ、ボルフィロモナス・ジンジバリス、スタフィロコッカス・アウレウス、コリネバクテリウム・ジフテリア、ノカルジア・コエリアカ、ストレプトコッカス・ニューモニアなどのバクテリア由来のDNAを具体的に挙げるができる。

【0017】かかる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質としては、非メチル化CpG配列を有する細菌DNAを特異的に認識することができるタンパク質であれば特に制限されるものではなく、例えば、配列表の配列番号2で示されるヒト由来のTLR9や、配列番号2で示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ上記非メチル化CpG配列を有する細菌DNAを特異的に認識することができるタンパク質や、これらの組換えタンパク質を具体的に挙げるができる。かかる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質は、そのDNA配列情報等に基づき公知の方法で調製することができる。

【0018】また、本発明の非メチル化CpG配列を有



する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAとしては、配列表の配列番号2で示されるヒト由来のTLR9をコードするDNA、例えば配列番号1で示されるものや、配列番号2で示されるアミノ酸配列において、1若しくは数個のアミノ酸が欠失、置換若しくは付加されたアミノ酸配列からなり、かつ上記非メチル化CpG配列を有する細菌DNAを特異的に認識することができるタンパク質をコードするDNAや、これらDNAとストリンジェントな条件下でハイブリダイズし、かつ上記非メチル化CpG配列を有する細菌DNAを特異的に認識することができるタンパク質をコードするDNAも包含され、これらはそのDNA配列情報等に基づき、例えばマウス由来のTLR9においてはマウスRAW264.7 cDNAライブラリーや129/SvJマウス遺伝子ライブラリーなどから公知の方法により調製することができる。

【0019】また、配列番号1に示される塩基配列又はその相補的配列並びにこれらの配列の一部又は全部をプローブとして、マウス由来のDNAライブラリーに対してストリンジェントな条件下でハイブリダイゼーションを行ない、該プローブにハイブリダイズするDNAを単離することにより、受容体タンパク質TLR9と同効な目的とする免疫誘導非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAを得ることもできる。かかるDNAを取得するためのハイブリダイゼーションの条件としては、例えば、42℃でのハイブリダイゼーション、及び1×SSC、0.1%のSDSを含む緩衝液による42℃での洗浄処理を挙げることができ、65℃でのハイブリダイゼーション、及び0.1×SSC、0.1%のSDSを含む緩衝液による65℃での洗浄処理をより好ましく挙げることができる。なお、ハイブリダイゼーションのストリンジェンシーに影響を与える要素としては、上記温度条件以外に種々の要素があり、当業者であれば、種々の要素を適宜組み合わせ、上記例示したハイブリダイゼーションのストリンジェンシーと同等のストリンジェンシーを実現することが可能である。

【0020】本発明の融合タンパク質とは、マウス、ヒト等の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に、マーカータンパク質及び/又はペプチドタグを結合させたものをいい、マーカータンパク質としては、従来知られているマーカータンパク質であればどのようなものでもよく、例えば、アルカリフォスファターゼ、抗体のFc領域、HRP、GFPなどを具体的に挙げることができ、また本発明におけるペプチドタグとしては、Mycタグ、Hisタグ、FLAGタグ、GSTタグなどの従来知られているペプチドタグを具体的に例示することができる。かかる融合タンパク質は、常法により作製することができ、NINTAとHisタグの親和性を利用した非メチル化Cp

G配列を有する細菌DNAを特異的に認識する受容体タンパク質の精製や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の検出や、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に対する抗体の定量や、その他当該分野の研究用試薬としても有用である。

【0021】本発明の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に特異的に結合する抗体としては、モノクローナル抗体、ポリクローナル抗体、キメラ抗体、一本鎖抗体、ヒト化抗体等の免疫特異的な抗体を具体的に挙げることができ、これらは上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を抗原として用いて常法により作製することができるが、その中でもモノクローナル抗体がその特異性の点でより好ましい。かかるモノクローナル抗体等の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に特異的に結合する抗体は、例えば、TLR9の変異又は欠失に起因する疾病の診断やTLR9の制御分子機構を明らかにする上で有用である。

【0022】非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に対する抗体は、慣用のプロトコルを用いて、動物（好ましくはヒト以外）に該非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質若しくはエピトープを含む断片、又は該タンパク質を膜表面に発現した細胞を投与することにより産生され、例えばモノクローナル抗体の調製には、連続細胞系の培養物により産生される抗体をもたらす、ハイブリドーマ法（Nature 256, 495-497, 1975）、トリオーマ法、ヒトB細胞ハイブリドーマ法（Immunology Today 4, 72, 1983）及びEBV-ハイブリドーマ法（MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp.77-96, Alan R. Liss, Inc., 1985）など任意の方法を用いることができる。以下に非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質として、マウス由来のTLR9を例に挙げてマウス由来のTLR9に対して特異的に結合するモノクローナル抗体、すなわち抗mTLR9モノクローナル抗体の作製方法を説明する。

【0023】上記抗mTLR9モノクローナル抗体は、抗mTLR9モノクローナル抗体産生ハイブリドーマをインビボ又はインビトロで常法により培養することにより生産することができる。例えば、インビボ系においては、齧歯動物、好ましくはマウス又はラットの腹腔内で培養することにより、またインビトロ系においては、動物細胞培養用培地で培養することにより得ることができる。インビトロ系でハイブリドーマを培養するための培地としては、ストレプトマイシンやペニシリン等の抗生物質を含むRPMI 1640又はMEM等の細胞培養培地を例示することができる。

【0024】抗mTLR9モノクローナル抗体産生ハイブリドーマは、例えば、マウス等から得られた受容体タンパク質TLR9を用いてBALB/cマウスを免疫し、免疫されたマウスの脾臓細胞とマウスNS-1細胞(ATCC TIB-18)とを、常法により細胞融合させ、免疫蛍光染色パターンによりスクリーニングすることにより、抗mTLR9モノクローナル抗体産生ハイブリドーマを作出することができる。また、かかるモノクローナル抗体の分離・精製方法としては、タンパク質の精製に一般的に用いられる方法であればどのような方法でもよく、アフィニティークロマトグラフィー等の液体クロマトグラフィーを具体的に例示することができる。

【0025】また、本発明の上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に対する一本鎖抗体をつくるためには、一本鎖抗体の調製法(米国特許第4,946,778号)を適用することができる。また、ヒト化抗体を発現させるために、トランスジェニックマウス又は他の哺乳動物等を利用したり、上記抗体を用いて、その非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現するクローンを単離・同定したり、アフィニティークロマトグラフィーでそのポリペプチドを精製することもできる。非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に対する抗体は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の分子機構を明らかにする上で有用である。

【0026】また上記抗mTLR9モノクローナル抗体等の抗体に、例えば、FITC(フルオレセインイソシアネート)又はテトラメチルローダミンイソシアネート等の蛍光物質や、<sup>125</sup>I、<sup>32</sup>P、<sup>35</sup>S又は<sup>3</sup>H等のラジオアイソトープや、アルカリホスファターゼ、ペルオキシダーゼ、β-ガラクトシダーゼ又はフィコエリトリン等の酵素で標識したものや、グリーン蛍光タンパク質(GFP)等の蛍光発光タンパク質などを融合させた融合タンパク質を用いることによって、上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の機能解析を行うことができる。また免疫学的測定方法としては、RIA法、ELISA法、蛍光抗体法、ブランク法、スポット法、血球凝集反応法、オクタロニー法等の方法を挙げることができる。

【0027】本発明はまた、上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現することができる発現系を含んでなる宿主細胞に関する。かかる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子の宿主細胞への導入は、Davisら(BASIC METHODS IN MOLECULAR BIOLOGY, 1986)及びSambrookら(MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold S

pring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989)などの多くの標準的な実験室マニュアルに記載される方法、例えば、リン酸カルシウムトランスフェクション、DEAE-デキストラン媒介トランスフェクション、トランスベクション(transvection)、マイクロインジェクション、カチオン性脂質媒介トランスフェクション、エレクトロポレーション、形質導入、スクレープローディング(scrape loading)、弾丸導入(ballistic introduction)、感染等により行うことができる。

そして、宿主細胞としては、大腸菌、ストレプトミセス、枯草菌、ストレプトコッカス、スタフィロコッカス等の細菌原核細胞や、酵母、アスペルギルス等の真菌細胞や、ドロソフィラS2、スポドプテラSf9等の昆虫細胞や、L細胞、CHO細胞、COS細胞、HeLa細胞、C127細胞、BALB/c3T3細胞(ジヒドロ葉酸レダクターゼやチミジンキナーゼなどを欠損した変異株を含む)、BHK21細胞、HEK293細胞、Bowesメラノーマ細胞、卵母細胞等の動植物細胞などを挙げることができる。

【0028】また、発現系としては、上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を宿主細胞内で発現させることができる発現系であればどのようなものでもよく、染色体、エピソーム及びウイルスに由来する発現系、例えば、細菌プラスミド由来、酵母プラスミド由来、SV40のようなパポバウイルス、ワクシニアウイルス、アデノウイルス、鶏痘ウイルス、仮性狂犬病ウイルス、レトロウイルス由来のベクター、バクテリオファージ由来、トランスポゾン由来及びこれらの組合せに由来するベクター、例えば、コスミドやファージミドのようなプラスミドとバクテリオファージの遺伝的要素に由来するものを挙げることができる。これら発現系は、発現を起こさせるだけでなく、発現を調節する制御配列を含んでいてもよい。

【0029】上記発現系を含んでなる宿主細胞やかかる細胞の細胞膜、またかかる細胞を培養して得られる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質は、後述するように本発明のスクリーニング方法に用いることができる。例えば、細胞膜を得る方法としては、F. Pietri-Rouxel (Eur. J. Biochem., 247, 1174-1179, 1997)らの方法などを用いることができ、また、かかる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を細胞培養物から回収し精製するには、硫酸アンモニウムまたはエタノール沈殿、酸抽出、アニオンまたはカチオン交換クロマトグラフィー、ホスホセルロースクロマトグラフィー、疎水性相互作用クロマトグラフィー、アフィニティークロマトグラフィー、ハイドロキシアパタイトクロマトグラフィーおよびレクチンクロマトグラフィーを含めた公知の方法、好ましくは、高速液体クロマトグラフィーが用いられる。特に、アフィニティークロマトグラ

フィーに用いるカラムとしては、例えば、抗 TLR9 モノクローナル抗体等の抗非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質抗体を結合させたカラムや、上記 TLR9 等の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質に通常のペプチドタグを付加した場合は、このペプチドタグに親和性のある物質を結合したカラムを用いることにより、これらの非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を得ることができる。

【0030】本発明において、上記非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現する非ヒト動物とは、野生型非ヒト動物に比べてかかる非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を大量に産生する非ヒト動物をいい、また、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物とは、染色体上の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子の一部若しくは全部が破壊・欠損・置換等の遺伝子変異により不活性化され、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質を発現する機能を失なった非ヒト動物をいう。そして、本発明における非ヒト動物としては、ウサギや、マウス、ラット等の齧歯目動物などの非ヒト動物を具体的に挙げることができるが、これらに限定されるものではない。

【0031】また、本発明において非メチル化 CpG 配列を有する細菌 DNA に対して不反応性とは、細菌 DNA による刺激に対する生体又は生体を構成する細胞、組織若しくは器官の反応性が低下しているか、あるいはほぼ失われていることを意味する。したがって、本発明において非メチル化 CpG 配列を有する細菌 DNA に対して不反応性の非ヒト動物とは、細菌 DNA による刺激に対して、生体又は生体を構成する細胞、組織若しくは器官の反応性が低下しているか、あるいはほぼ失われているマウス、ラット、ウサギ等のヒト以外の動物をいう。また、細菌 DNA による刺激としては、細菌 DNA を生体に投与するインビボでの刺激や、生体から分離された細胞に細菌 DNA を接触させるインビトロでの刺激等を挙げることができ、具体的には、TLR9 ノックアウトマウス等の TLR9 遺伝子機能が染色体上で欠損した非ヒト動物を挙げることができる。

【0032】ところで、メンデルの法則に従い出生してくるホモ接合体非ヒト動物には、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質欠損型又は過剰発現型とその同腹の野生型とが含まれ、これらホモ接合体非ヒト動物における欠損型又は過剰発現型とその同腹の野生型を同時に用いることによ

て個体レベルで正確な比較実験をすることができることから、野生型の非ヒト動物、すなわち非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損又は過剰発現する非ヒト動物と同種の動物、さらには同腹の動物を、例えば下記に記載する本発明のスクリーニングに際して併用することが好ましい。かかる非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損又は過剰発現する非ヒト動物の作製方法を、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質のノックアウトマウスやトランスジェニックマウスを例にとって以下説明する。

【0033】例えば、TLR9 等の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損したマウス、すなわち非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質ノックアウトマウスは、マウス遺伝子ライブラリーから PCR 等の方法により得られた遺伝子断片を用いて、上記非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子をスクリーニングし、スクリーニングされた非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子をウイルスベクター等を用いてサブクローンし、DNA シーケンシングにより特定する。このクローンの非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子の全部又は一部を pMC1 ネオ遺伝子カセット等に置換し、3' 末端側にジフテリアトキシン A フラグメント (DT-A) 遺伝子や単純ヘルペスウイルスのチミジンキナーゼ (HSV-tk) 遺伝子等の遺伝子を導入することによって、ターゲットベクターを作製する。

【0034】この作製されたターゲティングベクターを線状化し、エレクトロポレーション（電気穿孔）法等によって ES 細胞に導入し、相同的組換えを行い、その相同的組換え体の中から、G418 やガンシクロピア (GANC) 等の抗生物質により相同的組換えを起こした ES 細胞を選択する。また、この選択された ES 細胞が目的とする組換え体かどうかをサザンブロット法等により確認することが好ましい。その確認された ES 細胞のクローンをマウスの胚盤胞中にマイクロインジェクションし、かかる胚盤胞を仮親のマウスに戻し、キメラマウスを作製する。このキメラマウスを野生型のマウスとインタークロスさせると、ヘテロ接合体マウスを得ることができ、また、このヘテロ接合体マウスをインタークロスさせることによって、本発明の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質ノックアウトマウスを作製することができる。また、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識

する受容体タンパク質ノックアウトマウスが生起しているかどうかを確認する方法としては、例えば、上記の方法により得られたマウスからRNAを単離してノーザンブロット法等により調べたり、またこのマウスの発現をウエスタンブロット法等により調べる方法がある。

【0035】また、作出されたTLR9ノックアウトマウスが非メチル化CpG配列を有する細菌DNAに対して不応答性であることは、例えば、CpG ODNをTLR9ノックアウトマウスのマクロファージ、単核細胞、樹状細胞などの免疫細胞にインビトロ又はインビボで接触せしめ、かかる細胞におけるTNF- $\alpha$ 、IL-6、IL-12、IFN- $\gamma$ 等の産生量や、脾臓B細胞の増殖応答や、脾臓B細胞表面でのCD40、CD80、CD86、MHCクラスII等の抗原の発現量や、NF- $\kappa$ B、JNK、IRAK等のTLR9のシグナル伝達経路における分子の活性化を測定することにより確認することができる。そして、本発明のTLR9ノックアウトマウスは、非メチル化CpG配列を有する細菌DNA等の作用機序の解明や細菌感染に対するワクチン開発に有用なモデルとすることができる。

【0036】非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のトランスジェニックマウスは、TLR9等の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするcDNAにチキン $\beta$ -アクトリン、マウスニューロフィラメント、SV40等のプロモーター、及びラビット $\beta$ -グロビン、SV40等のポリA又はイントロンを融合させて導入遺伝子を構築し、該導入遺伝子をマウス受精卵の前核にマイクロインジェクションし、得られた卵細胞を培養した後、仮親のマウスの輸卵管に移植し、その後被移植動物を飼育し、産まれた仔マウスから前記cDNAを有する仔マウスを選択することによりかかるトランスジェニックマウスを創製することができる。また、cDNAを有する仔マウスの選択は、マウスの尻尾等より粗DNAを抽出し、導入した非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子をプローブとするドットハイブリダイゼーション法や、特異的プライマーを用いたPCR法等により行うことができる。

【0037】また、本発明の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAの全部あるいは一部を用いると、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の欠失又は異常に起因する疾病等の遺伝子治療に有効な細胞を調製することができる。本発明におけるこれら細胞の調製方法としては、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した細胞に、上記本発明のDNAの全部あるいは一部をトランスフェクション等により導入し、非メチル化Cp

G配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞を得る方法を挙げることができ、特に、かかる非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞としては、上記DNA等が染色体にインテグレートされ、ステイブルにTLR9活性を示す細胞を用いることが好ましい。

【0038】そしてまた、上記非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質とマーカータンパク質及び/又はペプチドタグとを結合させた融合非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質に対する抗体、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現することができる発現系を含んでなる宿主細胞、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現する非ヒト動物、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞等を用いると、本発明の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストや、非メチル化CpG配列を有する細菌DNAに対する反応性の抑制物質又は促進物質をスクリーニングすることができる。これらのスクリーニングにより得られたものは、細菌感染症に対する抑制物質又は促進物質や、アレルギー性疾患若しくは癌に対する抑制剤、予防剤又は治療薬や、遺伝子治療等において副作用を抑制剤又は阻害剤や、TLR9活性の欠失又は異常に起因する疾病等の診断・治療に有用な物質である可能性がある。

【0039】上記TLR9活性とは、非メチル化CpG配列を有する細菌DNAと特異的に反応し、細胞内にシグナルを伝達する機能をいい、シグナル伝達機能としては、TNF- $\alpha$ 、IL-6、IL-12、IFN- $\gamma$ 等のサイトカインを産生する機能や、亜硝酸イオンを産生する機能や、細胞を増殖する機能や、細胞表面においてCD40、CD80、CD86、MHCクラスII等の抗原を発現する機能や、NF- $\kappa$ B、JNK、IRAK等のTLR9のシグナル伝達経路における分子を活性化させる機能などを具体的に例示することができるが、これらに限定されるものではない。

【0040】本発明の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質のアゴニスト又はアンタゴニストのスクリーニング方法としては、被検物質の存在下、マクロファージ、脾臓細胞又は樹状細胞などの免疫細胞、非メチル化CpG配列を有す

る細菌DNAを特異的に認識する受容体タンパク質を発現している細胞、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質を発現する細胞等の非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質を発現している細胞をインビトロで培養し、TLR9活性を測定・評価する方法や、野生型非ヒト動物、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物、又は、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードする遺伝子が過剰発現した非ヒト動物に被検物質を投与し、該非ヒト動物から得られるマクロファージ、脾臓細胞、又は樹状細胞などの免疫細胞のTLR9活性を測定・評価する方法等を具体的に挙げることができる。

【0041】また、マクロファージ活性又は脾臓細胞活性の程度を測定・評価するに際し、対照として野生型非ヒト動物、特に同腹の野生型非ヒト動物の測定値と比較・評価することが個体差によるバラツキをなくすることができるので好ましい。このことは、以下に示す非メチル化CpG配列を有する細菌DNAに対する反応性の抑制物質又は促進物質のスクリーニングにおいても同様である。

【0042】また、非メチル化CpG配列を有する細菌DNAに対する反応性の抑制物質又は促進物質のスクリーニング方法としては、被検物質と非メチル化CpG配列を有する細菌DNAとの存在下、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質、又は該タンパク質を発現している細胞膜をインビトロでインキュベーションし、該タンパク質との反応性を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物から得られるマクロファージ又は脾臓細胞と被検物質とをあらかじめインビトロで接触せしめた後、該マクロファージ又は脾臓細胞を非メチル化CpG配列を有する細菌DNAの存在下で培養し、該マクロファージ若しくは脾臓細胞のマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物から得られるマクロファージ又は脾臓細胞と非メチル化CpG配列を有する細菌DNAとをあらかじめインビトロで接触せしめた後、該マクロファージ又は脾臓細胞を被検物質の存在下で培養し、該マクロファージ若しくは脾臓細胞のマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物にあらかじめ被検物質を投与した後、該非ヒト動物から得られるマクロ

ファージ又は脾臓細胞を非メチル化CpG配列を有する細菌DNAの存在下で培養し、該マクロファージ若しくは脾臓細胞のマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物にあらかじめ被検物質を投与した後、該非ヒト動物を細菌により感染させ、該非ヒト動物から得られるマクロファージ若しくは脾臓細胞のマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物をあらかじめ細菌により感染させた後、該非ヒト動物から得られるマクロファージ又は脾臓細胞を被検物質の存在下で培養し、該マクロファージ若しくは脾臓細胞のマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物にあらかじめ被検物質を投与した後、該非ヒト動物を細菌により感染させ、該非ヒト動物におけるマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法や、非メチル化CpG配列を有する細菌DNAに対して反応性を有するタンパク質をコードする遺伝子機能が染色体上で欠損した非ヒト動物をあらかじめ細菌により感染させた後、該非ヒト動物に被検物質を投与し、該非ヒト動物におけるマクロファージ活性又は脾臓細胞活性の程度を測定・評価する方法などを具体的に挙げることができる。また、これらのスクリーニング方法に用いる非メチル化CpG配列を有する細菌DNAとしては、CpG ODN (TCC-ATG-ACG-TTC-CTG-ATG-CT: 配列番号5) を用いることが好ましいが、これに限定されるものではない。

【0043】本発明はまた、検体中の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA配列を、本発明の非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNA配列と比較することからなる、非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質の活性又は発現と関連する疾病の診断に用いられる診断キットに関する。非メチル化CpG配列を有する細菌DNAを特異的に認識する受容体タンパク質をコードするDNAの変異型の検出は、遺伝子に変異がある個体をDNAレベルで見い出す

ことにより行うことができ、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質の過剰発現、過剰発現又は変異発現により生ずる疾病の診断に有効である。かかる検出に用いられる検体としては、被験者の細胞、例えば血液、尿、唾液、組織等の生検から得ることができるゲノム DNA や、RNA 又は cDNA を具体的に挙げるができるがこれらに限定されるものではなく、かかる検体を使用する場合、PCR 等により増幅したものをを用いることもできる。そして、塩基配列の欠失や挿入変異は、正常な遺伝子型と比較したときの増幅産物のサイズの変化により検出でき、また点突然変異は増幅 DNA を標識非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子とハイブリダイズさせることで同定することができる。このように、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする遺伝子の変異を検出することで、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質の活性又は発現と関連する疾病の診断又は判定をすることができる。

【0044】本発明はまた、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA 又は RNA のアンチセンス鎖の全部又は一部からなる非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質の活性又は発現と関連する疾患の診断用プローブ、及び当該プローブ及び／又は本発明の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質に特異的に結合する抗体を含有してなる非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質の活性又は発現と関連する疾患の診断キットに関する。前記診断用プローブとしては、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA (cDNA) 又は RNA (cRNA) のアンチセンス鎖の全部又は一部であり、プローブとして成立する程度の長さ (少なくとも 20 ベース以上) を有するものであれば特に制限されるものではない。かかるプローブ及び／又は本発明の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質に特異的に結合する抗体を細菌感染症等のような症状の疾患の診断薬の有効成分とするためには、プローブが分解されないような適当なバッファー類や滅菌水に溶解することが好ましい。また、これらの診断薬を用いた、免疫染色法 (Dev. Biol. 170, 207-222, 1995, J. Neurobiol. 29, 1-17, 1996) や、In situ ハイブリダイゼーション法 (J. Neurobiol. 29, 1-17, 1996) や、in situ PCR 法等の方法により細菌感染症等のような症状の疾患を診断することもできる。

【0045】本発明の医薬組成物としては、TLR9 等の非メチル化 CpG 配列を有する細菌 DNA を特異的に

認識する受容体タンパク質の全部又はその一部や、上記受容体タンパク質のアゴニストやアンタゴニストを含むものであれば、どのようなものでもよい。具体的には、細菌感染症に対するワクチンや、癌に対するワクチンや、気管支喘息をはじめとするアレルギー疾患の治療薬や、アンチセンスオリゴヌクレオチドを用いた治療や遺伝子治療において障害となる CpG モチーフの存在による副作用の克服剤・抑制剤・阻害剤などを挙げるができる。

【0046】前記のように、本発明の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA 配列の欠失、置換及び／又は付加に関連する疾病の診断キットとしては、TLR9 をコードする DNA を含むものであればどのようなものでもよく、かかる TLR9 をコードする DNA と検体中の非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA との塩基配列を比較することにより、非メチル化 CpG 配列を有する細菌 DNA を特異的に認識する受容体タンパク質をコードする DNA 配列の欠失、置換及び／又は付加に関連する疾病、例えば、癌、アレルギー、伝染病等の診断が可能となる。

【0047】

【実施例】以下に、実施例を挙げてこの発明を更に具体的に説明するが、この発明の技術的範囲はこれら実施例により限定されるものではない。

実施例 1 (TLR9 のクローニング)

ヒト TLR4 の DNA 配列情報を用いて、GenBank をサーチした結果、相同性がきわめて高いマウス EST (登録番号 AA273731; マウス) を見出した。このマウス EST の PCR 増幅産物をプローブとして、マウス RAW264.7 cDNA ライブラリーをスクリーニングし、完全な TLR9 オープンリーディングフレームを含む配列番号 3 に示される完全長の cDNA クローンを単離した。このマウス TLR9 の DNA 配列情報を用いて GenBank をサーチし、高い相同性を有するヒトゲノム配列を見出した。このヒトゲノム配列に基づいて、cDNA 端部を増幅し、U937 細胞 (J. Immunol. 163, 5039-5048, 1999) から、配列番号 1 に示される塩基配列を有する完全長のヒト TLR9 の cDNA を単離した。

【0048】実施例 2 (TLR9 ノックアウトマウスの作製)

129/SvJ マウス遺伝子ライブラリー (ストラタジーン社製) から TLR9 ゲノム DNA を単離し、pBluescript II SK(+) ベクター (ストラタジーン社製) 中でサブクローニングし、制限酵素マッピング及び DNA 配列決定により特定した。ターゲティングベクターは、TLR9 (ロイシンリッチリピート) 領域の一部分をコードする 1.0 kb のフラグメントを、ネオマイシン耐性遺伝子



カセット (pMCl-neo ; ストラタジーン社製) に置換し、負の選択マーカーとして単純ヘルペスウィルスチミジンキナーゼ (HSV-TK) を挿入することにより構築した (図 1)。このターゲティングベクターを線状化し、胎生 14. 1 日目の胚幹細胞 (ES 細胞) にエレクトポレーションし、G418 及びガンシクロビアに抵抗性を示す 292 個のクローンを選択し、PCR 法及びサザンブロット法により 14 個のクローンをスクリーニングした。

【0049】突然変異 TLR9 対立遺伝子を含有していた 3 個の標的 ES クローンを、C57BL/6 マウスの胚盤胞中にマイクロインジェクションしキメラマウスを作製した。この雄のキメラマウスを C57BL/6 雌マウスと交配させ、ヘテロ接合体 F1 マウスを作製し、かかるヘテロ接合体 F1 マウスをインタークロスすることによってホモ接合体マウス (TLR9 ノックアウトマウス: TLR9<sup>-/-</sup>) を得た (図 2)。なお、ホモ接合体マウスの確認は、マウスの尾から抽出した各ゲノム DNA を ScaI でダイジェストし、図 1 に示すプローブを用いたサザンブロット法により行った。本発明の TLR9 ノックアウトマウス (TLR9<sup>-/-</sup>) はメンデルの法則に従い作製することができ、12 週目までは顕著な異常を示さなかった。

【0050】突然変異により TLR9 遺伝子の不活性化が生起していることを確認するため、野生型マウス (+/+ ) 及び TLR9 ノックアウトマウス (-/-) の脾臓細胞から抽出した全 RNA (10 µg) を電気泳動にかけナイロン膜に移して、<sup>32</sup>P で標識した TLR9 の C-末端フラグメント若しくは N-末端フラグメント、又は β-アクチン (β-actin) に特異的な cDNA を用いてノーザンブロット分析を行った (図 3)。これらの結果から、TLR9 mRNA の N-末端フラグメントは TLR9 ノックアウトマウスの脾臓細胞からは検出されなかった。また、C-末端フラグメントをプローブとした場合、変異マウス由来の TLR9 の転写は野生型マウス由来のものとはほぼ同じサイズのものが検出されたが、生産量においては少ないことがわかった。そこで、変異マウスから得られた脾臓細胞の mRNA を用いて RT-PCR 法を行い、得られた生成物の配列分析を行った。この結果、転写された TLR9 遺伝子には neo 遺伝子が含まれており、この neo の挿入によって、TLR9 の N-末端部位にストップコドンが出現し、変異マウスにおいて機能的な TLR9 タンパク質が発現しないことがわかった (図 4)。なお、TLR9 ノックアウトマウスのリンパ細胞をフローサイトメトリーで測定した結果、異常成分は見られなかった。

【0051】実施例 3 (腹腔マクロファージの調製)  
野生型マウス (wild-type) 及び TLR9 ノックアウトマウス (TLR9<sup>-/-</sup>) のそれぞれの腹腔内に 4% のチオグリコール酸培地 (DIFCO 社製) を 2m

1 ずつ注入し、3 日後に各マウスの腹腔内から腹膜滲出細胞を単離し、これらの細胞を 10% のウシ胎仔血清 (GIBCO 社製) を添加した RPMI 1640 培地 (GIBCO 社製) 中で 37℃ にて 2 時間培養し、氷温のハンクス緩衝液 (Hank's buffered salt solution: HBSS; GIBCO 社製) で洗浄することにより非附着細胞を取り除き、附着細胞を腹膜マクロファージとして以下の実験に使用した。

【0052】実施例 4 (TLR9 ノックアウトマウスの非メチル化 CpG 配列を有する細菌 DNA に対する応答性)

最近、CpG ODN (oligodeoxynucleotide) の応答性は、TLR を介するシグナル伝達経路の中のアダプタータンパク質である MyD88 に依存していることが明らかになった。この MyD88 ノックアウトマウスは CpG ODN に対して応答しないが、TLR2 ノックアウトマウスや TLR4 ノックアウトマウスは正常に CpG ODN に対して応答する。これらのことは、CpG ODN が TLR2 及び TLR4 以外の TLR によって認識されることを示している。そこで、TLR9 ノックアウトマウスの CpG ODN に対する応答性を調べてみた。まず、腹腔マクロファージにおける炎症性サイトカインの産生量を以下のように測定した。

【0053】実施例 3 により調製した各腹膜マクロファージを INFγ (30 unit/ml) の存在下又は非存在下において、図 5 に示された各種濃度の CpG ODN (0. 1 又は 1. 0 µM; TIB MOLBIOL 社製; TC-C-ATG-ACG-TTC-CTG-ATG-CT)、PGN (10 µg/ml; Sigma and Fluka 社製; スタフィロコッカス・アウレウス由来)、LPS (1. 0 µg/ml; Sigma 社製; サルモネラ・ミネソタ Re-595 由来) といっしょに 24 時間培養した。培養後、培養上清中の TNFα、IL-6 及び IL-12 p40 の各濃度を ELISA 法により測定した。この結果を図 5 に示す。これらの結果から、野生型マウス (Wild-type) のマクロファージは CpG ODN に応答して TNFα、IL-6 及び IL-12 を産生し、さらに INFγ 及び CpG ODN で刺激すると、TNFα、IL-6 及び IL-12 の産生量が増加することがわかった。しかし、TLR9 ノックアウトマウス (TLR9<sup>-/-</sup>) 由来のマクロファージは、INFγ の存在下でさえ、CpG ODN に対する応答において検出可能なレベルの炎症性サイトカインを産生していなかった。また、野生型マウス及び TLR9 ノックアウトマウス由来のマクロファージは、LPS 又は PGN に対する応答により TNFα、IL-6 及び IL-12 をほぼ同程度産生することがわかった (図 5)。なお、それぞれの実験結果は n=3 の平均値を示す。図中の N.D. は検出できなかったことを示す。

【0054】また、CpG ODN 又は LPS に対する

野生型マウス (Wild-type) 及び TLR9 ノックアウトマウス (TLR9<sup>-/-</sup>) の脾臓細胞の応答性について調べてみた。それぞれのマウスの脾臓細胞 ( $1 \times 10^6$ ) を単離し、図 6 に示す各種濃度の CpG ODN 又は LPS により 96 ウェルプレート内で培養して脾臓細胞を刺激した。培養から 40 時間後に  $1 \mu\text{Ci}$  の [ $^3\text{H}$ ] - チミジン (デュポント社製) を添加して更に 8 時間培養し、[ $^3\text{H}$ ] の摂取量を  $\beta$  シンチレーションカウンター (パッカード社製) で測定した (図 6)。この結果から、野生型マウスの脾臓細胞では、CpG ODN や LPS の投与量に依存して細胞増殖反応を促進していたが、TLR9 ノックアウトマウスの脾臓細胞では、いかなる濃度の CpG ODN 刺激においても CpG ODN による細胞増殖反応は見られなかった。また、CpG ODN に応答して、野生型マウス由来の B 細胞表面の主要組織適合遺伝子複合体 (MHC) クラス II の発現が増加した。しかし、TLR9 ノックアウトマウス由来の B 細胞では CpG ODN に誘導された MHC クラス II の発現の増加は見られなかった。以上のことから、TLR9 ノックアウトマウスのマクロファージや B 細胞は、CpG ODN に対する応答性を特異的に欠如していることがわかった。

【0055】次に、CpG ODN を含有するバクテリア由来 DNA は樹状細胞を潜在的に刺激し、Th1 細胞の発達をサポートすることが知られている (EMBO J. 18, 6973-6982, 1999, J. Immunol. 161, 3042-3049, 1998, Proc. Natl. Acad. Sci. USA 96, 9305-9310, 1999)。そこで CpG ODN 誘導サイトカインの産生と、骨髄由来の樹状細胞の表面分子のアップレギュレーションを分析した。野生型マウス (Wild-type) 又は TLR9 ノックアウトマウス (TLR9<sup>-/-</sup>) の骨髄細胞を、 $10 \text{ ng/ml}$  のマウス顆粒球マクロファージコロニー刺激因子 (Peprotech 社製) を含む 10% のウシ胎仔血清を添加した RPMI 1640 培地で培養し (J. Exp. Med. 176, 1693-1702, 1992)、培養後 6 日目に未成熟の樹状細胞を回収し、 $0.1 \mu\text{M}$  の CpG ODN 又は  $0.1 \mu\text{g/ml}$  の LPS の存在下若しくは非存在下において、10% のウシ胎仔血清を添加した RPMI 1640 培地中で 2 日間培養した。培養後、上清中の IL-12 p40 の濃度を ELISA 法で測定した (図 7)。この結果から、野生型マウス由来の樹状細胞は CpG ODN に応答して IL-12 を産生したが、TLR9 ノックアウトマウス由来の樹状細胞においては、CpG ODN は IL-12 の産生を誘導しなかった。

【0056】上記  $10 \text{ ng/ml}$  のマウス顆粒球マクロファージコロニー刺激因子 (Peprotech 社製) を含む 10% のウシ胎仔血清を添加した RPMI 1640 培地で培養し、6 日目に回収された樹状細胞を、CD40、CD80、CD86 及び MHC クラス II に対する、それ

ぞれのビオチン化抗体により染色し、フィコエリトリン (phycoerythrin: PE; ファーミンジェン社製) で標識したストレプトアビジンで発展させ、これらの細胞をセルクレストソフトウェア (ベクトンディッキンソン社製) により蛍光活性化セルソーターキャリバー (FACS Calibur) で分析した (図 8)。この結果から、CpG ODN で刺激すると、野生型マウス由来の樹状細胞表面においては、CD40、CD80、CD86 及び MHC クラス II の発現を促進していたが、TLR9 ノックアウトマウス由来の樹状細胞表面では、CpG ODN に対する応答によりこれらの分子の発現を促進しなかった (図 8)。LPS による刺激では、野生型マウス由来の樹状細胞も TLR9 ノックアウトマウス由来の樹状細胞も同様の応答がみられた。以上の結果から、TLR9 は CpG ODN の細胞応答に不可欠な受容体であることがわかった。

【0057】実施例 5 (TLR9 ノックアウトマウス由来のマクロファージの CpG ODN に対する応答による NF- $\kappa$ B、JNK 及び IRAK の活性化) TLR のシグナルは、アダプター分子である MyD88 を介してセリン/トレオニンキナーゼである IRAK を活性化し、次いで MAP キナーゼ及び NF- $\kappa$ B を活性化することが知られている (Immunity 11, 115-122, 1999)。そこで CpG ODN が、かかる細胞内シグナル伝達分子を活性化するかどうかを調べてみた。実施例 3 により調製した野生型マウス及び TLR9 ノックアウトマウスの腹腔マクロファージ ( $1 \times 10^6$  cells) を、 $1.0 \mu\text{M}$  の CpG ODN 又は  $1.0 \mu\text{g/ml}$  のサルモネラ・ミネソタ Re-595 の LPS で図 9 に示された時間刺激し、各マウスのマクロファージから核蛋白質を抽出し、NF- $\kappa$ B の DNA 結合部位を含む特異的プローブと一時的にインキュベートし、電気泳動を行い、オートラジオグラフィーにより視覚化した (図 9)。

【0058】この結果から、CpG ODN で刺激すると、野生型マウス由来のマクロファージでは NF- $\kappa$ B の DNA 結合活性が増加するのに対し、TLR9 ノックアウトマウス由来のマクロファージでは NF- $\kappa$ B の DNA 結合活性は増加しなかった。TLR9 ノックアウトマウス由来のマクロファージを LPS で刺激したものは、野生型マウス由来のマクロファージを LPS で刺激したものと同様の NF- $\kappa$ B の活性化が見られた。以上の結果から、CpG ODN の誘導による NF- $\kappa$ B の活性が TLR9 ノックアウトマウス由来のマクロファージにおいて特異的に欠損していることがわかる。なお、図中の矢印は NF- $\kappa$ B と特異的プローブとの複合物の位置を示し、矢頭は特異的プローブのみの位置を示している。

【0059】上記と同様に図 10 又は図 11 で示された時間、CpG ODN 又は LPS で刺激した野生型マウ



ス及びTLR9ノックアウトマウスのマクロファージを、溶解緩衝液（最終濃度で1.0%のトリトンX-100、137mMのNaCl、20mMのトリス-HCl、5mMのEDTA、10%のグリセロール、1mMのPMSF、20 $\mu$ g/mlのアプロチニン、20 $\mu$ g/mlのロイペプチン、1mMのNa<sub>2</sub>VO<sub>4</sub>及び10mMの $\beta$ -グリセロリン酸を含有する緩衝液；pH8.

0) 中にて溶解し、この細胞溶解物を抗JNK抗体（サンタクルス社製）又は抗IRAK抗体（林原生化学研究所株式会社製）で免疫沈降して、文献（Immunity 11, 115-122, 1999）記載のように、インビトロキナーゼアッセイを行い、GST-c-Jun溶解蛋白質（GST-c-Jun）を基質としたJNK活性及びIRAKの活性を測定した（図10, 11における上段；GST-c-Jun, Auto）。

【0060】また、上記細胞溶解物を、SDS-ポリアクリルアミドゲル電気泳動により分離させ、ニトロセルロース膜に移し、この膜を抗JNK抗体（サンタクルス社製）又は抗IRAK抗体（Transduction Laboratories社製）でプロットして、エンハンスド・ケミルミネッセンス装置（デュボント社製）を使用して視覚化した（図10, 11における下段；WB）。以上の結果か

ら、CpG ODNは野生型マウス由来のマクロファージのJNK及びIRAKを活性化するが、TLR9ノックアウトマウス由来のマクロファージでは全く活性化しないことがわかった（図10, 11）。したがって、CpG ODNを介する情報伝達はTLR9に依存していることがわかった。

【0061】

【発明の効果】メチル化されていないCpGモチーフを含有するバクテリア由来DNAは免疫細胞を非常に活性化し、Th1の応答を誘導するが、そのバクテリア由来DNAを認識する受容体は知られていなかった。本発明により、細菌DNAの非メチル化CpG配列を含むオリゴヌクレオチドの受容体が明らかとなったことから、非メチル化CpG配列を有する細菌DNAを特異的に認識するTLRファミリーのメンバー受容体タンパク質TLR9や、それをコードする遺伝子DNA等は、細菌性疾病等の診断や、治療に用いることができ、またTLR9ノックアウト動物を用いると、バクテリア由来DNAの分子レベルにおける作用機作を明らかにすることが可能となる。

【0062】

【配列表】

#### SEQUENCE LISTING

<110> JAPAN SCIENCE AND TECHNOLOGY CORPORATION

<120> Specific receptor that recognizes bacterial DNA

<130> A031P63

<140>

<141>

<160> 5

<170> PatentIn Ver. 2.1

<210> 1

<211> 3257

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (107)..(3205)

<400> 1

ccgcctgcgc cccctggga agggaccctcg agtctgaagc atccctccct gtagctgctg 60  
tccagctgc cgcagacc ctctggagaa gccccgccc cccagc atg ggt ttc 115

Met Gly Phe

1

lgc cgc agc gcc ctg cac ccg ctg tct ctc ctg gtg cag gcc atc atg 163  
Cys Arg Ser Ala Leu His Pro Leu Ser Leu Leu Val Gln Ala Ile Met

5

10

15

ctg gcc atg acc ctg gcc ctg ggt acc ttg cct gcc ttc cta ccc tgt 211  
Leu Ala Met Thr Leu Ala Leu Gly Thr Leu Pro Ala Phe Leu Pro Cys

20

25

30

35

gag ctc cag ccc cac ggc ctg gtc aac tgc aac tgg ctg ttc ctg aag 259  
Glu Leu Gln Pro His Gly Leu Val Asn Cys Asn Trp Leu Phe Leu Lys

28

29	30
Gly Asn Leu Arg Val Leu Asp Leu Ser Glu Asn Phe Leu Tyr Lys Cys	
310	315 320
atc act aaa acc aag gcc ttc cag ggc cta aca cag ctg cgc aag ctt	1123
Ile Thr Lys Thr Lys Ala Phe Gln Gly Leu Thr Gln Leu Arg Lys Leu	
325	330 335
aac ctg tcc ttc aat tac caa aag agg gtg tcc ttt gcc cac ctg tct	1171
Asn Leu Ser Phe Asn Tyr Gln Lys Arg Val Ser Phe Ala His Leu Ser	
340	345 350 355
ctg gcc cct tcc ttc ggg agc ctg gtc gcc ctg aag gag ctg gac atg	1219
Leu Ala Pro Ser Phe Gly Ser Leu Val Ala Leu Lys Glu Leu Asp Met	
360	365 370
cac ggc atc ttc ttc cgc tca ctc gat gag acc acg ctc cgg cca ctg	1267
His Gly Ile Phe Phe Arg Ser Leu Asp Glu Thr Thr Leu Arg Pro Leu	
375	380 385
gcc cgc ctg ccc atg ctc cag act ctg cgt ctg cag atg aac ttc atc	1315
Ala Arg Leu Pro Met Leu Gln Thr Leu Arg Leu Gln Met Asn Phe Ile	
390	395 400
aac cag gcc cag ctc ggc atc ttc agg gcc ttc cct ggc ctg cgc tac	1363
Asn Gln Ala Gln Leu Gly Ile Phe Arg Ala Phe Pro Gly Leu Arg Tyr	
405	410 415
gtg gac ctg tcg gac aac cgc atc agc gga gct tcg gag ctg aca gcc	1411
Val Asp Leu Ser Asp Asn Arg Ile Ser Gly Ala Ser Glu Leu Thr Ala	
420	425 430 435
acc atg ggg gag gca gat gga ggg gag aag gtc tgg ctg cag cct ggg	1459
Thr Met Gly Glu Ala Asp Gly Gly Glu Lys Val Trp Leu Gln Pro Gly	
440	445 450
gac ctt gct ccg gcc cca gtg gac act ccc agc tct gaa gac ttc agg	1507
Asp Leu Ala Pro Ala Pro Val Asp Thr Pro Ser Ser Glu Asp Phe Arg	
455	460 465
ccc aac tgc agc acc ctc aac ttc acc ttg gat ctg tca cgg aac aac	1555
Pro Asn Cys Ser Thr Leu Asn Phe Thr Leu Asp Leu Ser Arg Asn Asn	
470	475 480
ctg gtg acc gtg cag ccg gag atg ttt gcc cag ctc tcg cac ctg cag	1603
Leu Val Thr Val Gln Pro Glu Met Phe Ala Gln Leu Ser His Leu Gln	
485	490 495
tgc ctg cgc ctg agc cac aac tgc atc tcg cag gca gtc aat ggc tcc	1651
Cys Leu Arg Leu Ser His Asn Cys Ile Ser Gln Ala Val Asn Gly Ser	
500	505 510 515
cag ttc ctg ccg ctg acc ggt ctg cag gtg cta gac ctg tcc cac aat	1699
Gln Phe Leu Pro Leu Thr Gly Leu Gln Val Leu Asp Leu Ser His Asn	
520	525 530
aag ctg gac ctc tac cac gag cac tca ttc acg gag cta cca cga ctg	1747
Lys Leu Asp Leu Tyr His Glu His Ser Phe Thr Glu Leu Pro Arg Leu	
535	540 545
gag gcc ctg gac ctc agc tac aac agc cag ccc ttt ggc atg cag ggc	1795
Glu Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe Gly Met Gln Gly	
550	555 560
gtg ggc cac aac ttc agc ttc gtg gct cac ctg cgc acc ctg cgc cac	1843
Val Gly His Asn Phe Ser Phe Val Ala His Leu Arg Thr Leu Arg His	
565	570 575

31	32
c t c a g c c t g g c c a c a a c a a c a t c c a c a g c c a a g t g t c c c a g c a g c t c	1891
Leu Ser Leu Ala His Asn Asn Ile His Ser Gln Val Ser Gln Gln Leu	
580	585 590 595
t g c a g t a c g t c g c t g c g g g c c c t g g a c t t c a g c g g c a a t g c a c t g g g c	1939
Cys Ser Thr Ser Leu Arg Ala Leu Asp Phe Ser Gly Asn Ala Leu Gly	
600	605 610
c a t a t g t g g g c c g a g g a g a c c t c t a t c t g c a c t t c t t c c a a g g c c t g	1987
His Met Trp Ala Glu Gly Asp Leu Tyr Leu His Phe Phe Gln Gly Leu	
615	620 625
a g c g g t t t g a t c t g g c t g g a c t t g t c c c a g a a c c g c c t g c a c a c c c t c	2035
Ser Gly Leu Ile Trp Leu Asp Leu Ser Gln Asn Arg Leu His Thr Leu	
630	635 640
c t g c c c c a a a c c c t g c g c a a c c t c c c c a a g a g c c t a c a g g t g c t g c g t	2083
Leu Pro Gln Thr Leu Arg Asn Leu Pro Lys Ser Leu Gln Val Leu Arg	
645	650 655
c t c c g t g a c a a t t a c c t g g c c t t c t t t a a g t g g t g g a g c c t c c a c t t c	2131
Leu Arg Asp Asn Tyr Leu Ala Phe Phe Lys Trp Trp Ser Leu His Phe	
660	665 670 675
c t g c c c a a a c t g g a a g t c t c g a c c t g g c a g g a a a c c a g c t g a a g g c c	2179
Leu Pro Lys Leu Glu Val Leu Asp Leu Ala Gly Asn Gln Leu Lys Ala	
680	685 690
c t g a c c a a t g g c a g c c t g c c t g c t g g c a c c c g g c t c c g g a g g c t g g a t	2227
Leu Thr Asn Gly Ser Leu Pro Ala Gly Thr Arg Leu Arg Arg Leu Asp	
695	700 705
g t c a g c t g c a a c a g c a t c a g c t t c g t g g c c c c g g c t t c t t t t c c a a g	2275
Val Ser Cys Asn Ser Ile Ser Phe Val Ala Pro Gly Phe Phe Ser Lys	
710	715 720
g c c a a g g a g c t g c g a g a g c t c a a c c t t a g c g c c a a c g c c c t c a a g a c a	2323
Ala Lys Glu Leu Arg Glu Leu Asn Leu Ser Ala Asn Ala Leu Lys Thr	
725	730 735
g t g g a c c a c t c c t g g t t t g g g c c c c t g g c g a g t g c c c t g c a a a t a c t a	2371
Val Asp His Ser Trp Phe Gly Pro Leu Ala Ser Ala Leu Gln Ile Leu	
740	745 750 755
g a t g t a a g c g c c a a c c c t c t g c a c t g c g c c t g t g g g g c g g c c t t t a t g	2419
Asp Val Ser Ala Asn Pro Leu His Cys Ala Cys Gly Ala Ala Phe Met	
760	765 770
g a c t t c c t g c t g g a g g t g c a g g c t g c c c g g t c t g c c c a g c c g g	2467
Asp Phe Leu Leu Glu Val Gln Ala Ala Val Pro Gly Leu Pro Ser Arg	
775	780 785
g t g a a g t g t g g c a g t c c g g g c c a g c t c c a g g g c c t c a g c a t c t t t g c a	2515
Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Leu Ser Ile Phe Ala	
790	795 800
c a g g a c c t g c g c c t c t g c c t g g a t g a g g c c c t c t c c t g g g a c t g t t t c	2563
Gln Asp Leu Arg Leu Cys Leu Asp Glu Ala Leu Ser Trp Asp Cys Phe	
805	810 815
g c c c t c t c g c t g c t g c t g t g g c c t g g g t g t g c c a t g c t g	2611
Ala Leu Ser Leu Leu Ala Val Ala Leu Gly Leu Gly Val Pro Met Leu	
820	825 830 835
c a t c a c c t c t g t g g c t g g g a c c t c t g g t a c t g c t t c c a c c t g t g c t g	2659
His His Leu Cys Gly Trp Asp Leu Trp Tyr Cys Phe His Leu Cys Leu	

33			34
	840	845	850
	gcc tgg ctt ccc tgg cgg ggg cgg caa agt ggg cga gat gag gat gcc		2707
	Ala Trp Leu Pro Trp Arg Gly Arg Gln Ser Gly Arg Asp Glu Asp Ala		
	855	860	865
	cig ccc tac gat gcc ttc gig gtc ttc gac aaa acg cag agc gca gtg		2755
	Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Thr Gln Ser Ala Val		
	870	875	880
	gca gac tgg gig tac aac gag ctt cgg ggg cag ctg gag gag tgc cgt		2803
	Ala Asp Trp Val Tyr Asn Glu Leu Arg Gly Gln Leu Glu Glu Cys Arg		
	885	890	895
	ggg cgc tgg gca ctc cgc ctg tgc ctg gag gaa cgc gac tgg ctg cct		2851
	Gly Arg Trp Ala Leu Arg Leu Cys Leu Glu Glu Arg Asp Trp Leu Pro		
	900	905	910
	ggc aaa acc ctc ttt gag aac ctg tgg gcc tgc gtc tat ggc agc cgc		2899
	Gly Lys Thr Leu Phe Glu Asn Leu Trp Ala Ser Val Tyr Gly Ser Arg		
	920	925	930
	aag acg ctg ttt gtg ctg gcc cac acg gac cgg gtc agt ggt ctc ttg		2947
	Lys Thr Leu Phe Val Leu Ala His Thr Asp Arg Val Ser Gly Leu Leu		
	935	940	945
	cgc gcc agc ttc ctg ctg gcc cag cag cgc ctg ctg gag gac cgc aag		2995
	Arg Ala Ser Phe Leu Leu Ala Gln Gln Arg Leu Leu Glu Asp Arg Lys		
	950	955	960
	gac gtc gtg gtg ctg gtg atc ctg agc cct gac ggc cgc cgc tcc cgc		3043
	Asp Val Val Val Leu Val Ile Leu Ser Pro Asp Gly Arg Arg Ser Arg		
	965	970	975
	tac gtg cgg ctg cgc cag cgc ctc tgc cgc cag agt gtc ctc ctc tgg		3091
	Tyr Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val Leu Leu Trp		
	980	985	990
	ccc cac cag ccc agt ggt cag cgc agc ttc tgg gcc cag ctg ggc atg		3139
	Pro His Gln Pro Ser Gly Gln Arg Ser Phe Trp Ala Gln Leu Gly Met		
	1000	1005	1010
	gcc ctg acc agg gac aac cac cac ttc tat aac cgg aac ttc tgc cag		3187
	Ala Leu Thr Arg Asp Asn His His Phe Tyr Asn Arg Asn Phe Cys Gln		
	1015	1020	1025
	gga ccc acg gcc gaa tag ccgtagccg gaatccigca cggigccacc		3235
	Gly Pro Thr Ala Glu		
	1030		
	tcacactca cctcaccctc gc		3257
	<210> 2		
	<211> 1032		
	<212> PRT		
	<213> Homo sapiens		
	<400> 2		
	Met Gly Phe Cys Arg Ser Ala Leu His Pro Leu Ser Leu Leu Val Gln		
	1 5 10 15		
	Ala Ile Met Leu Ala Met Thr Leu Ala Leu Gly Thr Leu Pro Ala Phe		
	20 25 30		
	Leu Pro Cys Glu Leu Gln Pro His Gly Leu Val Asn Cys Asn Trp Leu		
	35 40 45		
	Phe Leu Lys Ser Val Pro His Phe Ser Met Ala Ala Pro Arg Gly Asn		

36

37  
 450                      455                      460  
 Asp Phe Arg Pro Asn Cys Ser Thr Leu Asn Phe Thr Leu Asp Leu Ser  
 465                      470                      475                      480  
 Arg Asn Asn Leu Val Thr Val Gln Pro Glu Met Phe Ala Gln Leu Ser  
                     485                      490                      495  
 His Leu Gln Cys Leu Arg Leu Ser His Asn Cys Ile Ser Gln Ala Val  
                     500                      505                      510  
 Asn Gly Ser Gln Phe Leu Pro Leu Thr Gly Leu Gln Val Leu Asp Leu  
                     515                      520                      525  
 Ser His Asn Lys Leu Asp Leu Tyr His Glu His Ser Phe Thr Glu Leu  
                     530                      535                      540  
 Pro Arg Leu Glu Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe Gly  
 545                      550                      555                      560  
 Met Gln Gly Val Gly His Asn Phe Ser Phe Val Ala His Leu Arg Thr  
                     565                      570                      575  
 Leu Arg His Leu Ser Leu Ala His Asn Asn Ile His Ser Gln Val Ser  
                     580                      585                      590  
 Gln Gln Leu Cys Ser Thr Ser Leu Arg Ala Leu Asp Phe Ser Gly Asn  
                     595                      600                      605  
 Ala Leu Gly His Met Trp Ala Glu Gly Asp Leu Tyr Leu His Phe Phe  
                     610                      615                      620  
 Gln Gly Leu Ser Gly Leu Ile Trp Leu Asp Leu Ser Gln Asn Arg Leu  
 625                      630                      635                      640  
 His Thr Leu Leu Pro Gln Thr Leu Arg Asn Leu Pro Lys Ser Leu Gln  
                     645                      650                      655  
 Val Leu Arg Leu Arg Asp Asn Tyr Leu Ala Phe Phe Lys Trp Trp Ser  
                     660                      665                      670  
 Leu His Phe Leu Pro Lys Leu Glu Val Leu Asp Leu Ala Gly Asn Gln  
                     675                      680                      685  
 Leu Lys Ala Leu Thr Asn Gly Ser Leu Pro Ala Gly Thr Arg Leu Arg  
                     690                      695                      700  
 Arg Leu Asp Val Ser Cys Asn Ser Ile Ser Phe Val Ala Pro Gly Phe  
 705                      710                      715                      720  
 Phe Ser Lys Ala Lys Glu Leu Arg Glu Leu Asn Leu Ser Ala Asn Ala  
                     725                      730                      735  
 Leu Lys Thr Val Asp His Ser Trp Phe Gly Pro Leu Ala Ser Ala Leu  
                     740                      745                      750  
 Gln Ile Leu Asp Val Ser Ala Asn Pro Leu His Cys Ala Cys Gly Ala  
                     755                      760                      765  
 Ala Phe Met Asp Phe Leu Leu Glu Val Gln Ala Ala Val Pro Gly Leu  
                     770                      775                      780  
 Pro Ser Arg Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Leu Ser  
 785                      790                      795                      800  
 Ile Phe Ala Gln Asp Leu Arg Leu Cys Leu Asp Glu Ala Leu Ser Trp  
                     805                      810                      815  
 Asp Cys Phe Ala Leu Ser Leu Leu Ala Val Ala Leu Gly Leu Gly Val  
                     820                      825                      830  
 Pro Met Leu His His Leu Cys Gly Trp Asp Leu Trp Tyr Cys Phe His  
                     835                      840                      845  
 Leu Cys Leu Ala Trp Leu Pro Trp Arg Gly Arg Gln Ser Gly Arg Asp



39  
 850 855 860  
 Glu Asp Ala Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Thr Gln  
 865 870 875 880  
 Ser Ala Val Ala Asp Trp Val Tyr Asn Glu Leu Arg Gly Gln Leu Glu  
 885 890 895  
 Glu Cys Arg Gly Arg Trp Ala Leu Arg Leu Cys Leu Glu Glu Arg Asp  
 900 905 910  
 Trp Leu Pro Gly Lys Thr Leu Phe Glu Asn Leu Trp Ala Ser Val Tyr  
 915 920 925  
 Gly Ser Arg Lys Thr Leu Phe Val Leu Ala His Thr Asp Arg Val Ser  
 930 935 940  
 Gly Leu Leu Arg Ala Ser Phe Leu Leu Ala Gln Gln Arg Leu Leu Glu  
 945 950 955 960  
 Asp Arg Lys Asp Val Val Val Leu Val Ile Leu Ser Pro Asp Gly Arg  
 965 970 975  
 Arg Ser Arg Tyr Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val  
 980 985 990  
 Leu Leu Trp Pro His Gln Pro Ser Gly Gln Arg Ser Phe Trp Ala Gln  
 995 1000 1005  
 Leu Gly Met Ala Leu Thr Arg Asp Asn His His Phe Tyr Asn Arg Asn  
 1010 1015 1020  
 Phe Cys Gln Gly Pro Thr Ala Glu  
 1025 1030  
 <210> 3  
 <211> 3471  
 <212> DNA  
 <213> Mus musculus  
 <220>  
 <221> CDS  
 <222> (107)..(3205)  
 <400> 3  
 tgaagtgic acttccitcaa ttctctgaga gacccctggig tggaaatca ttctctgccg 60  
 cccagtttgt cagagggagc ctctgggagaa tcttccatct cccaac atg gtt ctc 115  
 Met Val Leu  
 1  
 cgt cga agg act ctg cac ccc ttg tcc ctc ctg gta cag gct gca glg 163  
 Arg Arg Arg Thr Leu His Pro Leu Ser Leu Leu Val Gln Ala Ala Val  
 5 10 15  
 ctg gct gag act ctg gcc ctg ggt acc ctg cct gcc ttc cta ccc tgt 211  
 Leu Ala Glu Thr Leu Ala Leu Gly Thr Leu Pro Ala Phe Leu Pro Cys  
 20 25 30 35  
 gag ctg aag cct cat ggc ctg glg gac tgc aat tgg ctg ttc ctg aag 259  
 Glu Leu Lys Pro His Gly Leu Val Asp Cys Asn Trp Leu Phe Leu Lys  
 40 45 50  
 tct gta ccc cgt ttc tct gcg gca gca tcc tgc tcc aac atc acc cgc 307  
 Ser Val Pro Arg Phe Ser Ala Ala Ala Ser Cys Ser Asn Ile Thr Arg  
 55 60 65  
 ctc tcc ttg atc tcc aac cgt atc cac cac ctg cac aac tcc gac ttc 355  
 Leu Ser Leu Ile Ser Asn Arg Ile His His Leu His Asn Ser Asp Phe  
 70 75 80

41	42
gtc cac ctg tcc aac ctg cgg cag ctg aac ctc aag tgg aac tgt cca	403
Val His Leu Ser Asn Leu Arg Gln Leu Asn Leu Lys Trp Asn Cys Pro	
85 90 95	
ccc act ggc ctt agc ccc ttg cac ttc tct tgc cac atg acc att gag	451
Pro Thr Gly Leu Ser Pro Leu His Phe Ser Cys His Met Thr Ile Glu	
100 105 110 115	
ccc aga acc ttc ctg gct atg cgt aca ctg gag gag ctg aac ctg agc	499
Pro Arg Thr Phe Leu Ala Met Arg Thr Leu Glu Glu Leu Asn Leu Ser	
120 125 130	
tat aat ggt atc acc act gtc ccc cga ctg ccc agc tcc ctg gtg aat	547
Tyr Asn Gly Ile Thr Thr Val Pro Arg Leu Pro Ser Ser Leu Val Asn	
135 140 145	
ctg agc ctg agc cac acc aac atc ctg gtt cta gat gct aac agc ctc	595
Leu Ser Leu Ser His Thr Asn Ile Leu Val Leu Asp Ala Asn Ser Leu	
150 155 160	
gcc ggc cta tac agc ctg cgc gtt ctc ttc atg gac ggg aac tgc tac	643
Ala Gly Leu Tyr Ser Leu Arg Val Leu Phe Met Asp Gly Asn Cys Tyr	
165 170 175	
tac aag aac ccc tgc aca gga gcg gtg aag gtg acc cca ggc gcc ctc	691
Tyr Lys Asn Pro Cys Thr Gly Ala Val Lys Val Thr Pro Gly Ala Leu	
180 185 190 195	
ctg ggc ctg agc aat ctc acc cat ctg tct gtg aag tat aac aac ctc	739
Leu Gly Leu Ser Asn Leu Thr His Leu Ser Val Lys Tyr Asn Asn Leu	
200 205 210	
aca aag gtg ccc cgc caa ctg ccc ccc agc ctg gag tac ctc ctg gtg	787
Thr Lys Val Pro Arg Gln Leu Pro Pro Ser Leu Glu Tyr Leu Leu Val	
215 220 225	
tcc tat aac ctc att gtc aag ctg ggg cct gaa gac ctg gcc aat ctg	835
Ser Tyr Asn Leu Ile Val Lys Leu Gly Pro Glu Asp Leu Ala Asn Leu	
230 235 240	
acc tcc ctt cga gta ctt gat gtg ggt ggg aat tgc cgt cgc tgc gac	883
Thr Ser Leu Arg Val Leu Asp Val Gly Gly Asn Cys Arg Arg Cys Asp	
245 250 255	
cat gcc ccc aat ccc tgt ata gaa tgt ggc caa aag tcc ctc cac ctg	931
His Ala Pro Asn Pro Cys Ile Glu Cys Gly Gln Lys Ser Leu His Leu	
260 265 270 275	
cac cct gag acc ttc cat cac ctg agc cat ctg gaa ggc ctg gtg ctg	979
His Pro Glu Thr Phe His His Leu Ser His Leu Glu Gly Leu Val Leu	
280 285 290	
aag gac agc tct ctc cat aca ctg aac tct tcc tgg ttc caa ggt ctg	1027
Lys Asp Ser Ser Leu His Thr Leu Asn Ser Ser Trp Phe Gln Gly Leu	
295 300 305	
gtc aac ctc tgc gtg ctg gac cta agc gag aac ttt ctc tat gaa agc	1075
Val Asn Leu Ser Val Leu Asp Leu Ser Glu Asn Phe Leu Tyr Glu Ser	
310 315 320	
atc aac cac acc aat gcc ttt cag aac cta acc cgc ctg cgc aag ctc	1123
Ile Asn His Thr Asn Ala Phe Gln Asn Leu Thr Arg Leu Arg Lys Leu	
325 330 335	
aac ctg tcc ttc aat tac cgc aag aag gta tcc ttt gcc cgc ctc cac	1171
Asn Leu Ser Phe Asn Tyr Arg Lys Lys Val Ser Phe Ala Arg Leu His	

43	44
340	355
cig gca agt tcc ttc aag aac ctg gtg tca ctg cag gag ctg aac atg	1219
Leu Ala Ser Ser Phe Lys Asn Leu Val Ser Leu Gln Glu Leu Asn Met	
360	370
aac ggc atc ttc ttc cgc tgc ctg aac aag tac acg ctg aga tgg ctg	1267
Asn Gly Ile Phe Phe Arg Ser Leu Asn Lys Tyr Thr Leu Arg Trp Leu	
375	385
gcc gat ctg ccc aaa ctg cac act ctg cat ctt caa atg aac ttc atc	1315
Ala Asp Leu Pro Lys Leu His Thr Leu His Leu Gln Met Asn Phe Ile	
390	400
aac cag gca cag ctg agc atc ttt ggt acc ttc cga gcc ctt cgc ttt	1363
Asn Gln Ala Gln Leu Ser Ile Phe Gly Thr Phe Arg Ala Leu Arg Phe	
405	415
gtg gac ttg tca gac aat cgc atc agt ggg cct tca acg ctg tca gaa	1411
Val Asp Leu Ser Asp Asn Arg Ile Ser Gly Pro Ser Thr Leu Ser Glu	
420	435
gcc acc cct gaa gag gca gat gat gca gag cag gag gag ctg ttg tct	1459
Ala Thr Pro Glu Glu Ala Asp Asp Ala Glu Gln Glu Glu Leu Leu Ser	
440	450
gcg gat cct cac cca gct cca ctg agc acc cct gct tct aag aac ttc	1507
Ala Asp Pro His Pro Ala Pro Leu Ser Thr Pro Ala Ser Lys Asn Phe	
455	465
atg gac agg tgt aag aac ttc aag ttc acc atg gac ctg tct cgg aac	1555
Met Asp Arg Cys Lys Asn Phe Lys Phe Thr Met Asp Leu Ser Arg Asn	
470	480
aac ctg gtg act atc aag cca gag atg ttt gtc aat ctg tca cgc ctg	1603
Asn Leu Val Thr Ile Lys Pro Glu Met Phe Val Asn Leu Ser Arg Leu	
485	495
cag tgt ctt agc ctg agc cac aac tcc att gca cag gct gtc aat ggc	1651
Gln Cys Leu Ser Leu Ser His Asn Ser Ile Ala Gln Ala Val Asn Gly	
500	515
tct cag ttc ctg ccg ctg act aat ctg cag gtg ctg gac ctg tcc cat	1699
Ser Gln Phe Leu Pro Leu Thr Asn Leu Gln Val Leu Asp Leu Ser His	
520	530
aac aaa ctg gac ttg tac cac tgg aaa tgc ttc agt gag cta cca cag	1747
Asn Lys Leu Asp Leu Tyr His Trp Lys Ser Phe Ser Glu Leu Pro Gln	
535	545
ttg cag gcc ctg gac ctg agc tac aac agc cag ccc ttt agc atg aag	1795
Leu Gln Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe Ser Met Lys	
550	560
ggt ala ggc cac aat ttc agt ttt gtg gcc cat ctg tcc atg cta cac	1843
Gly Ile Gly His Asn Phe Ser Phe Val Ala His Leu Ser Met Leu His	
565	575
agc ctt agc ctg gca cac aat gac att cat acc cgt gtg tcc tca cat	1891
Ser Leu Ser Leu Ala His Asn Asp Ile His Thr Arg Val Ser Ser His	
580	595
ctc aac agc aac tca gtg agg ttt ctt gac ttc agc ggc aac ggt atg	1939
Leu Asn Ser Asn Ser Val Arg Phe Leu Asp Phe Ser Gly Asn Gly Met	
600	610
ggc cgc atg tgg gat gag ggg ggc ctt tat ctg cat ttc ttc caa ggc	1987

45	46
Gly Arg Met Trp Asp Glu Gly Gly Leu Tyr Leu His Phe Phe Gln Gly	
615	620 625
ctg agt ggc ctg ctg aag ctg gac ctg tct caa aat aac ctg cat atc	2035
Leu Ser Gly Leu Leu Lys Leu Asp Leu Ser Gln Asn Asn Leu His Ile	
630	635 640
ctc cgg ccc cag aac ctt gac aac ctc ccc aag agc ctg aag ctg ctg	2083
Leu Arg Pro Gln Asn Leu Asp Asn Leu Pro Lys Ser Leu Lys Leu Leu	
645	650 655
agc ctc cga gac aac tac cta tct ttc ttt aac tgg acc agt ctg tcc	2131
Ser Leu Arg Asp Asn Tyr Leu Ser Phe Phe Asn Trp Thr Ser Leu Ser	
660	665 670 675
ttc ctg ccc aac ctg gaa gtc cta gac ctg gca ggc aac cag cta aag	2179
Phe Leu Pro Asn Leu Glu Val Leu Asp Leu Ala Gly Asn Gln Leu Lys	
680	685 690
gcc ctg acc aat ggc acc ctg cct aat ggc acc ctc ctc cag aaa ctg	2227
Ala Leu Thr Asn Gly Thr Leu Pro Asn Gly Thr Leu Leu Gln Lys Leu	
695	700 705
gat gtc agc agc aac agt atc gtc tct gtg gtc cca gcc ttc ttc gct	2275
Asp Val Ser Ser Asn Ser Ile Val Ser Val Val Pro Ala Phe Phe Ala	
710	715 720
ctg gcg gtc gag ctg aaa gag gtc aac ctc agc cac aac att ctc aag	2323
Leu Ala Val Glu Leu Lys Glu Val Asn Leu Ser His Asn Ile Leu Lys	
725	730 735
acg gtg gat cgc tcc tgg ttt ggg ccc att gtg atg aac ctg aca gtt	2371
Thr Val Asp Arg Ser Trp Phe Gly Pro Ile Val Met Asn Leu Thr Val	
740	745 750 755
cta gac gtg aga agc aac cct ctg cac tgt gcc tgt ggg gca gcc ttc	2419
Leu Asp Val Arg Ser Asn Pro Leu His Cys Ala Cys Gly Ala Ala Phe	
760	765 770
gta gac tta ctg ttg gag gtg cag acc aag gtg cct ggc ctg gct aat	2467
Val Asp Leu Leu Leu Glu Val Gln Thr Lys Val Pro Gly Leu Ala Asn	
775	780 785
ggt gtg aag tgt ggc agc ccc ggc cag ctg cag ggc cgt agc atc ttc	2515
Gly Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Arg Ser Ile Phe	
790	795 800
gca cag gac ctg cgg ctg tgc ctg gat gag gtc ctc tct tgg gac tgc	2563
Ala Gln Asp Leu Arg Leu Cys Leu Asp Glu Val Leu Ser Trp Asp Cys	
805	810 815
ttt ggc ctt tca ctc ttg gct gtg gcc gtg ggc atg gtg gtg cct ata	2611
Phe Gly Leu Ser Leu Leu Ala Val Ala Val Gly Met Val Val Pro Ile	
820	825 830 835
ctg cac cat ctc tgc ggc tgg gac gtc tgg tac tgt ttt cat ctg tgc	2659
Leu His His Leu Cys Gly Trp Asp Val Trp Tyr Cys Phe His Leu Cys	
840	845 850
ctg gca tgg cta cct ttg ctg gcc cgc agc cga cgc agc gcc caa gct	2707
Leu Ala Trp Leu Pro Leu Leu Ala Arg Ser Arg Arg Ser Ala Gln Ala	
855	860 865
ctc ccc tat gat gcc ttc gtg gtg ttc gat aag gca cag agc gca gtt	2755
Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Ala Gln Ser Ala Val	
870	875 880

47 48

gcg gac tgg gtg tat aac gag ctg cgg gtg cgg ctg gag gag cgg cgc 2803  
Ala Asp Trp Val Tyr Asn Glu Leu Arg Val Arg Leu Glu Glu Arg Arg  
885 890 895

ggt cgc cga gcc cta cgc tgg tgt ctg gag gac cga gat tgg ctg cct 2851  
Gly Arg Arg Ala Leu Arg Leu Cys Leu Glu Asp Arg Asp Trp Leu Pro  
900 905 910 915

ggc cag acg ctc ttc gag aac ctc tgg gct tcc atc tat ggg agc cgc 2899  
Gly Gln Thr Leu Phe Glu Asn Leu Trp Ala Ser Ile Tyr Gly Ser Arg  
920 925 930

aag act cta ttt gtg ctg gcc cac acg gac cgc gtc agt ggc ctc ctg 2947  
Lys Thr Leu Phe Val Leu Ala His Thr Asp Arg Val Ser Gly Leu Leu  
935 940 945

cgc acc agc ttc ctg ctg gct cag cag cgc ctg tgg gaa gac cgc aag 2995  
Arg Thr Ser Phe Leu Leu Ala Gln Gln Arg Leu Leu Glu Asp Arg Lys  
950 955 960

gac gtg gtg gtg tgg gtg atc ctg cgt ccg gat gcc cac cgc tcc cgc 3043  
Asp Val Val Val Leu Val Ile Leu Arg Pro Asp Ala His Arg Ser Arg  
965 970 975

tat gtg cga ctg cgc cag cgt ctc tgc cgc cag agt gtg ctc ttc tgg 3091  
Tyr Val Arg Leu Arg Gln Arg Leu Cys Arg Gln Ser Val Leu Phe Trp  
980 985 990 995

ccc cag cag ccc aac ggg cag ggg ggc ttc tgg gcc cag ctg agt aca 3139  
Pro Gln Gln Pro Asn Gly Gln Gly Gly Phe Trp Ala Gln Leu Ser Thr  
1000 1005 1010

gcc ctg act agg gac aac cgc cac ttc tat aac cag aac ttc tgc cgg 3187  
Ala Leu Thr Arg Asp Asn Arg His Phe Tyr Asn Gln Asn Phe Cys Arg  
1015 1020 1025

gga cct aca gca gaa tag ctcagagcaa cagctggaaa cagctgcatc 3235  
Gly Pro Thr Ala Glu  
1030

ttcagcccg gtccccgagt tgcctgcct gccctgcct gctttactac accgctattt 3295  
ggcaagtcg caatatatgc taccaagcca ccaggccac ggagcaaagg tggcagtaa 3355  
agggtagttt tcttccatg catctttcag gagagtgaag atagacacca gaccacaca 3415  
gaacaggact ggagttcatt cctgcccct ccacccact ttgcctgtct cigtat 3471

<210> 4  
<211> 1032  
<212> PRT  
<213> Mus musculus  
<400> 4

Met Val Leu Arg Arg Arg Thr Leu His Pro Leu Ser Leu Leu Val Gln  
1 5 10 15  
Ala Ala Val Leu Ala Glu Thr Leu Ala Leu Gly Thr Leu Pro Ala Phe  
20 25 30  
Leu Pro Cys Glu Leu Lys Pro His Gly Leu Val Asp Cys Asn Trp Leu  
35 40 45  
Phe Leu Lys Ser Val Pro Arg Phe Ser Ala Ala Ala Ser Cys Ser Asn  
50 55 60  
Ile Thr Arg Leu Ser Leu Ile Ser Asn Arg Ile His His Leu His Asn  
65 70 75 80  
Ser Asp Phe Val His Leu Ser Asn Leu Arg Gln Leu Asn Leu Lys Trp

49		85		90		95		50
Asn Cys Pro Pro Thr Gly Leu Ser Pro Leu His Phe Ser Cys His Met								
	100			105			110	
Thr Ile Glu Pro Arg Thr Phe Leu Ala Met Arg Thr Leu Glu Glu Leu								
	115			120			125	
Asn Leu Ser Tyr Asn Gly Ile Thr Thr Val Pro Arg Leu Pro Ser Ser								
	130			135			140	
Leu Val Asn Leu Ser Leu Ser His Thr Asn Ile Leu Val Leu Asp Ala								
145		150			155		160	
Asn Ser Leu Ala Gly Leu Tyr Ser Leu Arg Val Leu Phe Met Asp Gly								
	165			170			175	
Asn Cys Tyr Tyr Lys Asn Pro Cys Thr Gly Ala Val Lys Val Thr Pro								
	180			185			190	
Gly Ala Leu Leu Gly Leu Ser Asn Leu Thr His Leu Ser Val Lys Tyr								
	195			200			205	
Asn Asn Leu Thr Lys Val Pro Arg Gln Leu Pro Pro Ser Leu Glu Tyr								
210		215			220			
Leu Leu Val Ser Tyr Asn Leu Ile Val Lys Leu Gly Pro Glu Asp Leu								
225		230			235		240	
Ala Asn Leu Thr Ser Leu Arg Val Leu Asp Val Gly Gly Asn Cys Arg								
	245			250			255	
Arg Cys Asp His Ala Pro Asn Pro Cys Ile Glu Cys Gly Gln Lys Ser								
	260			265			270	
Leu His Leu His Pro Glu Thr Phe His His Leu Ser His Leu Glu Gly								
	275			280			285	
Leu Val Leu Lys Asp Ser Ser Leu His Thr Leu Asn Ser Ser Trp Phe								
	290			295			300	
Gln Gly Leu Val Asn Leu Ser Val Leu Asp Leu Ser Glu Asn Phe Leu								
305		310			315		320	
Tyr Glu Ser Ile Asn His Thr Asn Ala Phe Gln Asn Leu Thr Arg Leu								
	325			330			335	
Arg Lys Leu Asn Leu Ser Phe Asn Tyr Arg Lys Lys Val Ser Phe Ala								
	340			345			350	
Arg Leu His Leu Ala Ser Ser Phe Lys Asn Leu Val Ser Leu Gln Glu								
	355			360			365	
Leu Asn Met Asn Gly Ile Phe Phe Arg Ser Leu Asn Lys Tyr Thr Leu								
	370			375			380	
Arg Trp Leu Ala Asp Leu Pro Lys Leu His Thr Leu His Leu Gln Met								
385		390			395		400	
Asn Phe Ile Asn Gln Ala Gln Leu Ser Ile Phe Gly Thr Phe Arg Ala								
	405			410			415	
Leu Arg Phe Val Asp Leu Ser Asp Asn Arg Ile Ser Gly Pro Ser Thr								
	420			425			430	
Leu Ser Glu Ala Thr Pro Glu Glu Ala Asp Asp Ala Glu Gln Glu Glu								
	435			440			445	
Leu Leu Ser Ala Asp Pro His Pro Ala Pro Leu Ser Thr Pro Ala Ser								
	450			455			460	
Lys Asn Phe Met Asp Arg Cys Lys Asn Phe Lys Phe Thr Met Asp Leu								
465		470			475		480	
Ser Arg Asn Asn Leu Val Thr Ile Lys Pro Glu Met Phe Val Asn Leu								

51  
 485 490 495  
 Ser Arg Leu Gln Cys Leu Ser Leu Ser His Asn Ser Ile Ala Gln Ala  
 500 505 510  
 Val Asn Gly Ser Gln Phe Leu Pro Leu Thr Asn Leu Gln Val Leu Asp  
 515 520 525  
 Leu Ser His Asn Lys Leu Asp Leu Tyr His Trp Lys Ser Phe Ser Glu  
 530 535 540  
 Leu Pro Gln Leu Gln Ala Leu Asp Leu Ser Tyr Asn Ser Gln Pro Phe  
 545 550 555 560  
 Ser Met Lys Gly Ile Gly His Asn Phe Ser Phe Val Ala His Leu Ser  
 565 570 575  
 Met Leu His Ser Leu Ser Leu Ala His Asn Asp Ile His Thr Arg Val  
 580 585 590  
 Ser Ser His Leu Asn Ser Asn Ser Val Arg Phe Leu Asp Phe Ser Gly  
 595 600 605  
 Asn Gly Met Gly Arg Met Trp Asp Glu Gly Gly Leu Tyr Leu His Phe  
 610 615 620  
 Phe Gln Gly Leu Ser Gly Leu Leu Lys Leu Asp Leu Ser Gln Asn Asn  
 625 630 635 640  
 Leu His Ile Leu Arg Pro Gln Asn Leu Asp Asn Leu Pro Lys Ser Leu  
 645 650 655  
 Lys Leu Leu Ser Leu Arg Asp Asn Tyr Leu Ser Phe Phe Asn Trp Thr  
 660 665 670  
 Ser Leu Ser Phe Leu Pro Asn Leu Glu Val Leu Asp Leu Ala Gly Asn  
 675 680 685  
 Gln Leu Lys Ala Leu Thr Asn Gly Thr Leu Pro Asn Gly Thr Leu Leu  
 690 695 700  
 Gln Lys Leu Asp Val Ser Ser Asn Ser Ile Val Ser Val Val Pro Ala  
 705 710 715 720  
 Phe Phe Ala Leu Ala Val Glu Leu Lys Glu Val Asn Leu Ser His Asn  
 725 730 735  
 Ile Leu Lys Thr Val Asp Arg Ser Trp Phe Gly Pro Ile Val Met Asn  
 740 745 750  
 Leu Thr Val Leu Asp Val Arg Ser Asn Pro Leu His Cys Ala Cys Gly  
 755 760 765  
 Ala Ala Phe Val Asp Leu Leu Leu Glu Val Gln Thr Lys Val Pro Gly  
 770 775 780  
 Leu Ala Asn Gly Val Lys Cys Gly Ser Pro Gly Gln Leu Gln Gly Arg  
 785 790 795 800  
 Ser Ile Phe Ala Gln Asp Leu Arg Leu Cys Leu Asp Glu Val Leu Ser  
 805 810 815  
 Trp Asp Cys Phe Gly Leu Ser Leu Leu Ala Val Ala Val Gly Met Val  
 820 825 830  
 Val Pro Ile Leu His His Leu Cys Gly Trp Asp Val Trp Tyr Cys Phe  
 835 840 845  
 His Leu Cys Leu Ala Trp Leu Pro Leu Leu Ala Arg Ser Arg Arg Ser  
 850 855 860  
 Ala Gln Ala Leu Pro Tyr Asp Ala Phe Val Val Phe Asp Lys Ala Gln  
 865 870 875 880  
 Ser Ala Val Ala Asp Trp Val Tyr Asn Glu Leu Arg Val Arg Leu Glu



<400> 5  
tccatgacgt tccatgatgt

20

【図面の簡単な説明】

【図 1】本発明の TLR9 ノックアウトマウスと野生型マウスの遺伝子地図を示す図である。

【図2】本発明のTLR9ノックアウトマウスのサザン  
ブロット分析の結果を示す図である。

【図3】本発明のTLR9ノックアウトマウスの脾臓細胞におけるノーザンブロット分析の結果を示す図である。

【図 4】本発明の TLR9 ノックアウトマウスと野生型マウスのアミノ酸配列の比較結果を示す図である。

【図5】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN、PGN又はLPS誘導によるTNF $\alpha$ 、IL-6又はIL12の産生量の結果を示す図である。

【図6】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導による細胞増殖応答の結果を示す図である。

【図7】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導によるIL-12の産生量の結果を示す図である。


【図8】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導によるCD40、CD80、CD86及びMHCクラスIIの発現量の結果を示す図である。

【図9】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導によるNF- $\kappa$ Bの活性化の結果を示す図である。


【図１０】本発明のTLR 9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導によるJNKの活性化の結果を示す図である。

【図 11】本発明のTLR9ノックアウトマウス及び野生型マウスにおけるCpG ODN又はLPS誘導によるIRAKの活性化の結果を示す図である。


【図 1】




【図 2】




【図 3】




【図 4】




【図 5】




【図 6】




【図 7】




【図 8】




【図 9】



【図10】



【図11】



フロントページの続き

(51) Int. Cl. <sup>7</sup>	識別記号	F I	テマコード (参考)
A 6 1 P 37/08		C 0 7 K 16/28	4 C 0 8 4
C 0 7 K 14/005		C 1 2 N 1/15	4 H 0 4 5
	16/28		1/19
C 1 2 N 1/15			1/21
	1/19	C 1 2 Q 1/68	A
	1/21	G 0 1 N 33/15	Z
	5/10		33/50
C 1 2 Q 1/68			33/566
G 0 1 N 33/15			33/577
	33/50	C 1 2 P 21/08	B
	33/566	(C 1 2 Q 1/68	A
	33/577	C 1 2 R 1:91)	
// C 1 2 P 21/08		(C 1 2 P 21/08	
(C 1 2 Q 1/68		C 1 2 R 1:91)	
C 1 2 R 1:91)		C 1 2 N 15/00	Z N A A
(C 1 2 P 21/08		A 6 1 K 37/02	
C 1 2 R 1:91)		C 1 2 N 5/00	B

Fターム(参考) 2G045 AA34 AA35 BB01 BB20 BB24  
BB51 CB01 CB17 DA36 FB02  
FB03 FB05 FB08  
4B024 BA44 BA63 CA04 DA02 DA06  
DA12 EA04 FA10 GA05 HA14  
HA15  
4B063 QA01 QA19 QQ43 QR08 QR32  
QR42 QR56 QS25 QS34 QX05  
4B064 AG27 CA10 CA20 CC24 DA15  
4B065 AA26X AA58X AA72X AA90Y  
AA91X AA92X AB01 AB05  
AC10 BA02 CA24 CA25 CA43  
4C084 AA01 AA17 BA44 ZB132  
ZB262 ZB352  
4H045 AA10 AA11 AA20 AA30 BA10  
CA40 DA50 EA50 EA52 FA72  
FA74

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**